

*A bit more on
Frequency-domain EEG*

and then...

*The Event-related Brain Potential
(Part 1)*

**Synchronization and
Desynchronization**

- Supposition that alpha blocking meant that the EEG had become desynchronized
 - Yet the activity is still highly synchronized -- not at 8-13 Hz
 - May involve fewer neuronal ensembles in synchrony



**Event-related
Synchronization and Desynchronization**

- Pfurtscheller (1992) -- Two types of ERS
 - Secondary (follows ERD)

Announcements


- Papers: 1 or 2 paragraph prospectus due a few minutes ago!
 - Feedback coming soon if you've not received it already
- 3x5s

If Alpha Desynchs, what Synchs?





Event-related Synchronization and Desynchronization

- Pfurtscheller (1992) -- Two types of ERS
 - Secondary (follows ERD)
 - Primary (Figure 3 & Figure 4)



Alpha power time course during reading (upper) and voluntary finger movements (lower). Primary ERS is seen over electrodes overlying cortical areas not involved in the task.



Primary ERS seen over parietal and occipital leads during right finger movement. ERD is seen over central electrodes, with earlier onset over hemisphere contralateral to movement.

Frontal Midline Theta (more later in advanced topics)

- Increased midline frontal theta during periods of high cognitive demand
- This is specifically under conditions in which cortical resources must be allocated for select cognitive processes
 - Attention
 - Memory
 - Error Monitoring

Saueng Hoppe Klimesch Gerloff Hummel (2007)

- Complex finger movement sequences
- Varied Task Difficulty, and Memory Load (2x2 design)
- Task-related Theta Power (4-7 Hz) computed for each condition relative to 5 min. resting baseline
- Phase coherence also examined across sites
 - Phase Locking Value (0-1)
 - Then expressed as percent increase over rest

Theta Power





FIG. 1. Task-related theta (4-7 Hz) power increase. White indicates a strong task-related power increase compared with rest. Note that only during execution of novel and complex sequences is strong frontal-midline theta exhibited. This indicates that frontal theta activity reflects both memory load and sequence complexity.

Saueng Hoppe Klimesch Gerloff Hummel (2007)

40 Hz Activity


- First reports of important 40 Hz activity
- Sheer & Grandstaff (1969) review
 - pronounced rhythmic electrical bursting
- Daniel Sheer's subsequent work until his death renewed interest in "40 Hz" phenomena



Note specificity of response to S_D , over visual cortex to discriminative stimulus, in 40-Hz range; Some hint of it later in the motor cortex. Note also decreased activity in slower bands during the same time periods.

Sheer work with Cats

- Learning paradigm
- Cat must learn
 - press to S_D (7cps light flicker)
 - not $S-$ (3 cps light flicker)
 - the hypothesis is that the synchronized 40 Hz activity represents the focused activation of specific cortical areas necessary for performance of a task



Note very different pattern to $S-$. No 40-Hz change in visual cortex, and marked increase in lower frequencies at same time period.


Human Studies

- Hypothesis is that 40 Hz activity correlates with the behavioral state of focused arousal (Sheer, 1976) or cortical activation
 - a "circumscribed state of cortical excitability" (Sheer, 1975)
 - Bird et al (1978)
 - biofeedback paradigm
 - increased 40 Hz activity is associated with high arousal and mental concentration
 - Ford et al., (1980)
 - subjects once trained to voluntarily suppress 40 Hz EEG are unable to maintain that suppression while simultaneously solving problems
 - concluded that problem solving and absence of 40 Hz are incompatible

Lateralized Task Effects

- Loring & Sheer (1984)
 - right-handed students
 - analogies task
 - spatial Task
- Results transformed into laterality ratios:
 - $(L-R)/(L+R)$ 40 Hz
 - higher # => greater LH activity (P3-O1-T5 triangle vs P4-O2-T6 triangle);
- Results
 - greatest variability during baseline
 - smallest variability and greatest LH activation during verbal
 - no laterality effects in the 40Hz EMG bands

Laterality of 40 Hz



Controlling for EMG contributions

- Spydell & Sheer (1982)
 - used similar tasks and found similar results
 - using conservative controls for muscle artifact

Hy, 1982 Alpha, Beta II, 40 Hz EEG, and 40 Hz EMG Activity 423

TABLE 1
Median changes in rate scores

Problems	Alpha		Beta II		40 Hz Total		40 Hz EMG		40 Hz EMG	
	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right
Verbal	-36.7*	-52.4*	-20.1*	-20.2*	1.0*	0.1	1.2*	0.1	8.4*	10.6*
Reaction	-36.1*	-37.6*	-15.3*	-15.3*	0.7	1.0*	0.4	0.9*	13.9*	8.5*


*p < .05.

Spydell and Sheer Vol.

TABLE 3
Spearman rank-order correlations between various 40 Hz activity measures

40 Hz Measure	Correlations			
	Verbal Left	Verbal Right	Notational Left	Rotational Right
40 Hz EEG	.74*	.68*	.94*	.78*
40 Hz EMG	.27	.28	.25	.25

*p < .05.




Individual Differences

- Spydell & Sheer (1983), Alzheimers
 - controls showed task related changes in EEG with appropriate lateralization
 - Alz did not
- Schnyer & Allen (1995)
 - Most highly hypnotizable subjects showed enhanced 40 hz activity

So this is exciting, why didn't this work take off immediately?

- The EMG concern
 - The concern is likely over-rated (recall Table 3)
- Sheer died
- But not all is lost, as there is renewed interest...



Mukamel et al *Science* 2005
 recorded single unit activity and local field potentials in auditory cortex of two neurosurgical patients and compared them with the fMRI signals of 11 healthy subjects during presentation of an identical movie segment. The predicted fMRI signals derived from single units and the measured fMRI signals from auditory cortex showed a highly significant correlation.

Singer (1993)

- Revitalized interest in the field

The Binding Problem


- Potentially infinite number of things and ideas that we may attempt to represent within the CNS
 - Cells code for limited sets of features,
 - These must somehow be integrated
 - -- the so-called binding problem
- If there exists a cell for a unique contribution of attributes, then convergent information from many cells could converge on such a cell
 - But there are a finite # of cells and interconnections
- And even the billions and billions of cells we have cannot conceivably handle the diversity of representations

The Functional Perspective -- as yet merely a theory

- There is no site of integration
 - Integration is achieved through simultaneous activation of an assembly of neurons distributed across a wide variety of cortical areas
 - Neurons in such assemblies must be able to adaptively identify with other neurons within the assembly while remaining distinct from other neurons in other assemblies
 - This association with other neurons is through a temporal code of firing (Synchronicity)
 - This even allows for the possibility that a single neuron could be part of two active assemblies (via a multitasking procedure)


Implications

- Also allows for the possibility that there exists no direct neuronal connection between neurons within an assembly
 - merely the fact that they are simultaneously activated that makes the unified experience of the object possible
- Yet what can synchronize these oscillations?



Jensen et al, *TICS*, 2012

Implications – Alpha as a synchronization mechanism



Jensen et al, *TICS*, 2012

Functional Role of Gamma Synchronization

- Feedforward coincidence detection
 - To summate effectively, signals must arrive at post-synaptic neuron from multiple sources within msec of each other (else decay)
 - Gamma-band synchronization can lead to temporal focusing of inputs from multiple and distributed pre-synaptic neurons
- Rhythmic Input Gain Modulation
 - Excitatory input is most effective when it arrives out of phase with inhibitory input and vice versa
 - Allows for precision and efficiency of signal transmission (or inhibition)

Fries, 2009

Implications

- This view is a dynamic view
 - depends on experience
 - can change with experience
- Synchronously activated units more likely to become enhanced and part of an assembly that will subsequently become synchronously activated
- Singer concludes:
 - Points out the problem of looking for synchronous activation on the micro level, suggesting that a return to the EEG literature looking for task-dependent synchronization in the gamma (aka 40 Hz) band!
- Forty-Hz may indeed make a comeback!
 - "Forty" = $40 \pm$ some range
 - Gamma! (Stay tuned during advanced topics)

The Event-Related Potential (aka the ERP)

Overview

Event-related potentials are patterned voltage changes embedded in the ongoing EEG that reflect a process in response to a particular event: e.g., a visual or auditory stimulus, a response, an internal event






Figure 4.2. A schematic representation of ERP components elicited by auditory, infrequent target stimuli. The three panels represent three different voltage \times time functions: the left bottom panel shows the very early sensory components (with a latency of less than 10 ms); the left top panel shows the middle latency sensory components (with a latency of between 10 and 50 ms); and the right panel shows late components (latency exceeding 50 ms). Note the different voltage and time scales used in the three panels, as well as the different nomenclatures used to label the peaks (components). (Adapted with permission of the author from Donchin, 1979, with kind permission of Springer Science and Business media.)

Time-locked activity
and extraction by
averaging



The Classic View:

Time-locked activity and extraction by signal averaging

- Ongoing activity reflects "noise"
- Activity that reflects processing of a given stimulus "signal"
- The signal-related activity can be extracted because it is **time-locked** to the presentation of the stimulus
- Signal Averaging is most common method of extracting the signal
 - Sample EEG for ~1 second after each stimulus presentation & average together across like stimuli
 - Time-locked signal emerges; noise averages to zero
 - Signal to noise ratio increases as a function of the **square root** of the number of trials in the average

What does the ERP reflect?

- May reflect sensory, motor, **and/or** cognitive events in the brain
- Reflect the synchronous and phase-locked activities of large neuronal populations engaged in information processing

Component is a "bump" or "trough"




Figure 4.2. A schematic representation of ERP components elicited by auditory, infrequent target stimuli. The three panels represent three different voltage \times time functions: the left bottom panel shows the very early sensory components (with a latency of less than 10 ms); the left top panel shows the middle latency sensory components (with a latency of between 10 and 50 ms); and the right panel shows late components (latency exceeding 50 ms). Note the different voltage and time scales used in the three panels, as well as the different nomenclatures used to label the peaks (components). (Adapted with permission of the author from Donchin, 1979, with kind permission of Springer Science and Business media.)

Making Meaning from the bumps

Pores o'er the Cranial map with learned eyes,
Each rising hill and bumpy knoll decries
Here secret fires, and there deep mines of sense
His touch detects beneath each prominence.



Nomenclature & Quantifying

- Most commonly label peaks and troughs by polarity (P or N) and latency at active recording site
- Quantifying
 - Amplitude
 - Latency
 - Area
 - "String" measure
 - Fancy stuff to be discussed in "advanced" topics

Component is a "bump" or "trough"




Figure 4.2. A schematic representation of ERP components elicited by auditory, infrequent target stimuli. The three panels represent three different voltage \times time functions: the left bottom panel shows the very early sensory components (with a latency of less than 10 ms); the left top panel shows the middle-latency sensory components (with a latency of between 10 and 50 ms); and the right panel shows late components (latency exceeding 50 ms). Note the different voltage and time scales used in the three panels, as well as the different nomenclatures used to label the peaks (components). (Adapted with permission of the author from Donchin, 1979, with kind permission of Springer Science and Business media.)

Early Components

- Waves I-VI represent evoked activity in auditory pathways and nuclei of the brainstem
- Early components <60-100 msec
 - occur in **obligatory** fashion
 - are called **Exogenous** = determined "outside" organism
- Even subtle deviations in appearance may be indicative of pathology

Later ERP components

- Highly sensitive to changes in
 - State of organism
 - Meaning of stimulus (NOT physical characteristics)
 - Information processing demands of task
- Therefore termed **Endogenous** = determined "within" organism

Not all components fit neatly into exogenous or endogenous categories

- Both Obligatory but modulated by psychological factors
- "Mesogenous"

Defining Components:


aka how do I know one when I see one?

- By positive and negative peaks at various latencies and scalp locations
- By functional associations, covarying across subjects, conditions, or scalp locations in response to experimental manipulations
- By neuronal structures that plausibly give rise to them

Evoked Vs Emitted ERP's


- Evoked are most commonly studied: occur in response to a physical stimulus
- Emitted potentials occur in absence of a physical stimulus (e.g., omission of item in sequence)
- Evoked can have both exogenous and endogenous components; emitted usually have only endogenous

After Fabiani, Gratton, Federmeier, 2007




Comparison to other "windows on the brain"

- Very precise temporal resolution



Comparison to other "windows on the brain"

- Very precise temporal resolution
- Spatial localization is more difficult
 - At the surface, activity of many functional synaptic units recorded
 - ERP's generated only by groups of cells that are synchronously activated in a geometrically organized manner



After Lorente de N6, 1947




Figure 1-11. Anatomy and electrogenesis of ventroposterior (VP) thalamus. A. Horizontal section showing bushy arborizations of lemniscal (lem) afferents terminating on dendrites of VP relay neurons (r). (From "Patterns of Organization in Specific and Non-specific Thalamic Fields" by M. E. Scheibel and A. B. Scheibel. In D. P. Purpura and M. D. Yahr [Eds.], *The Thalamus*. New York: Columbia University Press, 1966. Reprinted by permission.) B. Postulated potential field produced by depolarization of VP relay neurons. For clarity, the most intense parts of the field are omitted.

Comparison to other "windows on the brain"

- Very precise temporal resolution
- Spatial localization is more difficult
 - At the surface, activity of many functional synaptic units recorded
 - ERP's generated only by groups of cells that are synchronously activated in a geometrically organized manner
 - Synchronous activation may occur in one or more than one location
 - Monopolar recording technique most often used
 - Yet localization is not impossible in conjunction with other techniques

Correlate Vs substrate (AGAIN)

- Late ERP components should not be taken to indicate the existence of a neurological substrate of cognitive processing
- Rather should be considered a correlate
- Constructs in search of validation; Process of validation:
 - Determine antecedent conditions under which the ERP component appears and also magnitude and latency of ERP component
 - Develop hypotheses concerning functional significance of the "subroutine" underlying the ERP component
 - Predict consequences of subroutine--validate empirically

Basic Signal Processing

Caveat Emptor

- DO NOT interpret scalp distribution of ERP's as reflect cortical specialization
- Also, DO NOT interpret area of maximum amplitude to suggest that generator lies underneath




Paradigms and acquisition

- Precise temporal control over stimulus presentation necessary
 - Requires discrete stimuli or responses
- Individual stimuli are presented numerous times; ERP's generally do not habituate, unlike peripheral measures
- Concurrent with each stimulus, a signal/pulse must be sent to the A/D converter to indicate time of stimulus onset
- Sampling epochs (legacy!) vs continuously
 - Considerations for sampling epochs
 - pre-onset samples (to provide a baseline for comparison)
 - epoch length
- Epochs for like stimuli averaged together to create ERP for that set of stimuli

Assumptions of Averaging methods


- Signal and noise (in each epoch) sum linearly together to produce the recorded waveform for each epoch (not some peculiar interaction)
- The evoked signal waveshape attributable **solely** to the stimulus is the same for each presentation
- The noise contributions can be considered to constitute statistically independent samples of a random process

Demo of Averaging




Filtering and its influence on the ERP

- Despite many trials and averaging, some noise may remain in the averaged waveform
- If you are only interested in later & slower components, then a low-pass filter may be of interest



Same ERP filtered with 12.5 (black), 8 (red), and 5 (lime) Hz Low Pass FIR Filter




Same ERPs overlaid; note amplitude attenuation in P3 amplitude with stricter filters


Let's ERP!

Applications of Early Components

- Neurological evaluation of sensory function; e.g. evaluation of hearing in infants
- Tones of various dB intensities presented and V wave in auditory brainstem ERP examined
- **Figure 10**; 4000 individual trials per average



Prediction of recovery from coma




- ❑ Somatosensory evoked potentials were recorded from a patient who was still comatose 1 week after severe closed head injury.
- ❑ Responses evoked by electrical stimulation of left and right median nerves
- ❑ Normal tracing seen at Erb's point, and from the next over vertebra prominens, but not over C3 of C4.
- ❑ Absence of any cortical response a bad prognostic sign. Patient continued in a chronic vegetative state 1 year after accident

Inter-Hemispheric Transfer Time (IHTT)

- Hypothesized that interhemispheric transfer of information may be abnormal in various disorders (e.g., dyslexia)
- Reaction Time measures contain too much variability not related to Transfer Time
- ERP early components appear promising as a measure of time required to transfer information between hemispheres

IHTT Study (Saron)

- Checkerboards subtending < 1 degree of visual angle presented 2.9 degrees from center
- ERP's recorded at O1 and O2
- Problem of lateralization and Paradoxical results possible: parafoveal regions on banks of calcarine fissure
- P100 wave latency examined; earlier latency in occiput contralateral to presentation
 - Measured by peak picking procedure
 - Also by cross-lagged correlation technique
 - Both methods suggest ~15 millisecond IHTT; found to be in expected direction predicted by anatomy for over 90% of subjects
 - Reaction time data from same task showed no reliable differences



P1, N1, and Attention





Fig. 1. Paradigm for using ERPs to study attention. Stimulus display (left) and idealized results (right). Subjects fixate a central cross and attend either to the left or right visual field. Stimuli are then presented to the left and right visual fields in a rapid sequence. In this example, the ERP elicited by a left visual field stimulus contains larger P1 and N1 components when the stimulus is attended ('Attend left') than when it is ignored ('Attend right').

From Luck et al, *TICS*, 2000

More than Spatial Directed Attention



Increases stimulus complexity results in more rapid early processing

Note:
Amplitude of P1
Latency of P1
Latency of N1

Taylor
Clinical Neurophys
2002

Fig. 2. Grand averaged visual ERPs at Pz electrode for the 3 array sizes, showing the shorter latencies, larger P1s for array size 17, but longer latency P3 (dark arrows) than for array sizes 5 and 9 (grey arrows). These are averaged across colour, orientation and conjunction conditions, as this ERP effect was seen regardless of whether it was a single feature or conjunction trial.

More than Spatial Directed Attention





Fig. 3. Mean P1 latencies across 7 age groups, showing the consistently shorter latencies to faces compared to inverted faces and control stimuli (phase-scrambled faces and flowers). There were 15 children in each of the 6 age groups and 38 adults (adapted from Taylor et al., 2001c).

Taylor
Clinical Neurophys
2002

“These combined PET/ERP data therefore provide strong evidence that sustained visual spatial attention results in a preset, top-down biasing of the early sensory input channels in a retinotopically organized way”



Woldorff et al., *Human Brain Mapping*, 1997

Prelude to Advance Topic: Source Localization

Observed Potentials

Model Potentials Dorsal Occipital PET Seeds





Figure 3. Left: Observed potential distributions in the attend-left-minus-attend-right difference waves at the peak of the P1 attention effect (110–130 msec). Right: Corresponding model potential distributions seeded by the dorsal occipital PET foci, which provided an excellent fit to the P1 effect (residual variance 2%).

P1 and Sleep

P1 REAPPEARANCE DURING REM

SUBJECT K.K.



Note P1 disappears in Stage 2 sleep, but reemerges in REM sleep

Construct Validity of P300 (P3, P3b)

- First observed by Sutton, Braren, Zubin, & John (1965)
- P300 Amplitude; Johnson's model is

$$P300 \text{ Amplitude} = f[T \times (1/P + M)]$$
 where
 - P = probability of occurrence,
 - M = Stimulus meaning, &
 - T = amount of information transmitted





Figure 2. Grand-mean waveforms (N=7) from Fz, Cz, and Pz from three different tasks. The ERPs elicited in an oddball paradigm run under two different task conditions, Counting (solid line) and Reaction Time (dashed line), are superimposed on the ERP elicited when the same stimulus signified correct performance in a feedback paradigm (dotted line). The waveforms were all elicited by a 1000 Hz, 50dB SL tone ($p=50$).

Aspects of the Model

- Rarity
 - The P300 is observed in variants of the "oddball paradigm"
 - The rare stimulus almost invariably elicits a P300: largest at parietal, then central, and then frontal sites
 - Subjective probability
- Stimulus meaning
 - Actually composed of three dimensions
 - task complexity
 - stimulus complexity
 - stimulus value
- Information Transmission (proportion 0 to 1; example)




Figure 12-1. The ERPs in each column were elicited by the same physical tone; high-pitched tones were used for the left column and low-pitched tones for the right column. Both were presented in a Bernoulli series in which the probability of the two stimuli were equal. In the middle of each column (labeled "A") is the ERP elicited by all the presentations of the stimulus. The curve labeled "AA" was obtained by averaging together all the tones of one frequency that were preceded on the previous trial by tones of the same frequency. On the other hand, the curves labeled "BA" were elicited by stimuli preceded on the previous trial by the tones of different frequency. Similar sorting operations were applied to all other curves in this figure. It can be seen that the same physical tone elicited quite different ERPs, depending on the events that occurred on the preceding trials. Whenever a tone terminated a series of tones from the other category, a large P300 was elicited, and its magnitude was a function of the length of the stimulus series. (From "Effect of Stimulus Sequence on the Waveform of the Cortical Event-Related Potential," by K. C. Squires, C. D. Wickens, N. K. Squires, and E. Donchin. Science, 1976, 193, 1142-1146. Copyright 1976 by the AAAS.

Information Transmission




Fig. 2. Grand averaged visual ERPs at Pz electrode for the 3 array sizes, showing the shorter latencies, larger P1s for array size 17, but longer latency P3 (dark arrows) than for array sizes 5 and 9 (grey arrows). These are averaged across colour, orientation and conjunction conditions, as this ERP effect was seen regardless of whether it was a single feature or conjunction trial.

Taylor
Clinical Neurophys
2002

P3 Latency

- An index of processing time, independent of response requirements
 - RT measures confounds the two
 - McCarthy & Donchin (1981) experiment:
 - The words "RIGHT" or "LEFT" embedded in a matrix of letters of X's
 - Compatible condition: respond with hand indicated in matrix; Incompatible condition: respond with opposite hand (e.g., LEFT signals right hand response);
 - Results:
 - P300 latency delayed when discriminability more difficult
 - Response compatibility had no effect on P300 latency
 - Note amplitude reduction as function of noise-information transmission)





Figure 4.10. ERP waveforms at Pz averaged across subjects for three different semantic categorization tasks. The solid line indicates ERPs obtained during a task in which the subjects had to distinguish between the word DAVID and the word NANCY (the FN condition). The dotted line indicates ERPs obtained during a task in which the subjects had to decide whether a word presented was a male or a female name (the VN condition). The dashed line indicates ERPs obtained during a task in which the subjects had to decide whether a word was or was not a synonym of the word PROD (the SYN condition). These three tasks were considered to involve progressively more difficult discriminations. Note the latency of P300 peak is progressively longer as the discrimination is made more difficult. (Copyright 1977, AAAS. Adapted with permission of the author and publisher from Kutas, McCarthy, & Donchin, 1977.)

Not only difficulty in physical discrimination, but difficulty in cognitive categorization


Construct Validity?

- What, then, does the P300 mean in very general terms?
 - A stimulus (or class of stimuli) is "important"; denotes information that is necessary or useful to the task
 - Stimulus is meaningful, important, noticeable
 - Evaluated within context of working memory? (cf. Donchin & Coles, 1988; Verlager 1988; Polich, 2007; Verlager, 2008)
- The P3a (Squires, Squires, and Hillyard, 1975): P3-like component with a frontal maximum and occurs to improbable stimuli in the "to-be-ignored" class of stimuli; a novelty response.

How Many P3s?



- The Classic P3/P300
 - Parietal Central Maximum
 - Largest when stimuli rare and task-relevant
- The P3a (Squires et al., 1975) or Novelty P3 (Courchesne et al., 1975)
 - More anterior scalp distribution
 - Slightly earlier latency
 - Responsive to rare, unexpected, unattended stimuli



Simons et. al, 2001

- Squires Task was tones (two tones)
- Courchesne task was digitized speech (“me” “you” and collection of naturally occurring sounds)
- In all cases subjects merely counted Tones

P3a – Can you see it?

- Some inconsistencies in finding P3a following the initial Squires, Squires and Hilyard 1975 report
- Comerchero & Polich (1998) may have resolved the enigma
 - P3a highly dependent on foreground discrimination





Fig. 1. ERP waveforms (left) and PCA basis waves (right) obtained from infrequent targets during the Squires (top) task and infrequent nontargets/levels during the Courchesne (bottom) task. PCA was conducted during the 230–420 ms epoch following stimulus onset and four factors were extracted from each data set.

Modality	Auditory		Visual	
	Low	High	Low	High
Nontarget distinctiveness				
Target (0.10)	2000 Hz 75 dB	2000 Hz 75 dB	12.57 cm ² ● Blue	12.57 cm ² ● Blue
Standard (0.80)	1940 Hz 75 dB	1940 Hz 75 dB	10.18 cm ² ● Blue	10.18 cm ² ● Blue
Nontarget (0.10)	500 Hz 75 dB	4000 Hz 90 dB	12.57 cm ² ■ Blue	12.57 cm ² ■ Fuchsia



Note: Nontarget peak amplitude was earlier and larger at the frontal electrodes than those from the target stimuli, but especially when foreground discrimination is difficult

Comerchero & Polich (1998), *Clinical Neurophysiology*




Fig. 1. Schematic illustration of the single-stimulus (top), oddball (middle), and three-stimulus (bottom) paradigms, with the elicited ERPs from the stimuli of each task at the right (Polich and Crafoio, 2006). The single-stimulus task presents an infrequent target (T) in the absence of any other stimuli. The oddball task presents two different stimuli in a random sequence, with one occurring less frequently than the other (target = T, standard = S). The three-stimulus task is similar to the oddball with a compelling distractor (D) stimulus that occurs infrequently. In each task, the subject is instructed to respond only to the target and otherwise to refrain from responding. The distractor elicits a P3a, and target elicits a P3b (P300). Reprinted with permission of the authors and from Elsevier (Copyright 2006).

Polich, *Clin Neurophys*, 2007

Synopsis

“...the manipulation of target-standard stimulus discriminability produced a stimulus environment in which the infrequently occurring nontarget engaged focal attention in a manner similar to that observed previously for ‘novel’ stimuli.”

“However, all stimuli in the present study were employed because of their ‘typical’ characteristics, so that the results imply that an anterior P3a component can be produced without using ‘novel’ stimuli per se.”

“If stimulus context is defined primarily by a difficult target/standard discrimination, attentional redirection to the nontarget would occur because of the frontal lobe activation that generates P3a.”

Comerchero & Polich 1998, p. 47

ERPs and Memory

- Sensitive to both Recognition
 - Likely episodic recollection
- Sensitive to Encoding

Repetition Priming Effects

- Robust effect that repeated items produce an enhanced late positivity across a broad latency range
- Magnitude of effect related to strength of memory trace






Fig. 4. Grand mean ERP waveforms elicited by correctly recognized old and correctly rejected new items from Johnson et al. (1998a). The left column depicts the old and new waveforms at the electrode site and hemisphere where that subcomponent was largest. Reproduced from Johnson et al. (1998a) with permission of the publisher.

Repetition Priming

- Are there repetition effects that do not depend on the subjective awareness of the subject?
 - Can use Masked Priming to examine (Schnyer, Allen, Forster, 1997)




Schnyer, Allen, Forster, 1997




Standard Repetition Effect for Words Seen Unmasked in Previous Blocks
Task is to make OLD-NEW decision

Schnyer, Allen, Forster, 1997




Standard Repetition Effect for Words Seen Unmasked in Previous Blocks But Task is to make WORD-NONWORD decision
Schnyer, Allen, Forster, 1997



Masked Repetition Priming Effect for Words Presented only a Trial Previously
Schnyer, Allen, Forster, 1997

Memory Encoding

- Words subsequently remembered show enhanced positivity at encoding
- Strategy interacts, however



Note prototypic DM effect on left, but not on right for those that used elaborative strategies. Note enhancement over frontal lead for these latter subjects.

Figure 4-12. ERPs elicited by "isolated" words that were later recalled (solid line) or not-recalled (dashed line). The left column shows ERPs for subjects who used rote mnemonic strategies; the right column shows ERPs for subjects who used elaborative strategies. Note that the amplitude of P300 is related to subsequent recall for the rote memorizers, but not for elaborators. (Copyright 1986, Elsevier Science Publishers. Reprinted with permission of the publisher from Fabiani, Karis, & Donchin, 1986b.)




Fig. 3. A: Grand mean ERPs elicited by study items that were subsequently associated with remember or know judgments (hits) or were unrecognized (misses) during the subsequent recognition test. B: Grand mean difference waveforms computed by subtracting the ERPs to study items subsequently missed from those that were subsequently associated with either a remember or know judgment (Modified from Friedman and Trott, 2000). C: CSD maps for 2 intervals (500–800, 810–1100 ms) measured in the Dm waveform associated with a subsequent Remember judgment. Data in A and B recorded at a left inferior prefrontal scalp site.

Indirect Assessments of Recognition

- Can the ERP detect recognition, independent of subjects' overt responses?
- Two applications
 - Clinical Malingering
 - Forensic Assessment

ERP Memory Assessment Procedures


- Learn a list of words
- Learn a second list of words
- Task: Concealed (1st list) and Nonconcealed (2nd list) words appear infrequently

Item Type	Probability	Response	P3 Amplitude
Nonconcealed	1/7	"Yes"	Large
Concealed	1/7	"No"	Large if Recognized Small if not Recognized
Unlearned	5/7	"No"	Small

- Similar to procedures by Rosenfeld et al, Farwell & Donchin

Motivational Variations


Conceal	Lie	Lie + \$\$
➤ "YES" for words <u>JUST</u> learned, "NO" for all others	➤ "YES" for words learned	➤ "YES" for words learned
➤ Try to hide the fact that you learned the first list of words I taught you	➤ Lie about words from the first list I taught you	➤ Lie about words from the first list I taught you
		➤ \$5.00 incentive



After Allen & Iacono, 1997

The Challenge

To provide statistically supported decisions for each and every subject, despite considerable individual variability in ERP morphology



Bayesian Combination of ERP Indicators: Probability that an ERP was elicited by Learned Items

Subject	List						
	Learned		Unlearned				
	NonConceal	Conceal	U1	U2	U3	U4	U5
#01	1.0	0.999	0.000	0.000	0.000	0.000	0.001
#02	1.0	1.0	0.000	0.000	0.000	0.000	0.000
#03	1.0	0.999	0.000	0.000	0.000	0.002	0.000
#04	1.0	1.0	0.000	0.001	0.002	0.000	0.000
#05	1.0	0.971	0.002	0.000	0.000	0.000	0.000
#06	1.0	0.999	0.000	0.000	0.000	0.000	0.000
#07	0.983	1.0	0.000	0.000	0.000	0.000	0.000
...							
#18	0.996	0.983	0.874	0.001	0.000	0.000	0.000
#19	0.009	0.214	0.971	0.000	0.002	0.189	0.983
#20	1.0	0.999	0.002	0.000	0.009	0.000	0.214

Note: Only trials in which subjects did not acknowledge concealed items included

Classification Accuracy based on ERPs

	Learned (true pos)	Unlearned (true neg)
Conceal	0.95	0.96
Lie	0.93	0.94
Lie + \$\$	0.95	0.98
Combined	0.94	0.96

Allen, Iacono, & Danielson, *Psychophysiology*, 1992

Brain Fingerprinting: A New Paradigm in Criminal Investigations and Counterterrorism

Executive Summary

Farwell Brain Fingerprinting is a revolutionary new technology for investigating crimes and exonerating innocent suspects, with a record of 100% accuracy in research on FBI agents, research with US government agencies, and field applications.

The technology is proprietary and patented. Brain Fingerprinting fulfills an urgent need for government, law enforcement agencies, corporations, and individuals. Over a trillion dollars are spent annually on crime fighting worldwide.

Brain Fingerprinting solves the central problem by determining scientifically whether a suspect has the details of a crime stored in his brain. It has received extensive media coverage around the world. The technology is fully developed and available for application.

Brain Fingerprinting is a powerful tool for the investigation of suspected terrorists. Measuring the brain wave activity while suspects are shown words or pictures related to specifics of the September 11, 2001 attacks can help determine if they are members of terrorist cells. Brain Fingerprinting can identify trained terrorists before they strike



Rev. Jerry Farwell

Larry Farwell, PhD
Chairman & Chief Scientist
Brain Wave Science
Human Brain Research Laboratory, Inc.

www.brainwavescience.com

Brain Fingerprinting Laboratories
a new paradigm....

Criminal Justice | Medical | Advertising | Security Testing

In the News | Research | Contact Us

Counterterrorism Applications

How do we determine if a person is a terrorist or spy? There is a new technology, that for the first time, allows us to measure scientifically if specific information is stored in a person's brain. Brain Fingerprinting technology can determine the presence or absence of specific information, such as terrorist training and associations. This exciting new technology can help address the following critical elements in the fight against terrorism:

- Aid in determining who has participated in terrorist acts, directly or indirectly.
- Aid in identifying trained terrorists with the potential to commit future terrorist acts, even if they are in a "sleeper" cell and have not been active for years.
- Help to identify people who have knowledge or training in banking, finance or communications and who are associated with terrorist teams and acts.
- Help to determine if an individual is in a leadership role within a terrorist organization.




The Claim


- Brain Fingerprinting can determine “scientifically whether a suspect has details of a crime stored in his brain”
- Thus these ERP-procedures should be able to identify memories in laboratory studies
- Two tests of the robustness of this procedure:
 - False recollections
 - Virtual Reality Mock Crime

A Laboratory Paradigm for False Recollections: DRM


- Subjects presented with 15 words highly associated with an omitted critical item



Reported Rates of Recognition



Allen and Mertens (in press)



Allen and Mertens (in press)

The Box Score Blues

Ground Truth	Test Verdict
	Recognized
Actually Learned	56%
Critical Lure	72%
Unlearned	4%

- Highlights the need to have memorable items in the test
- Suggests limited utility in substantiating disputed memories; e.g., claims regarding recovered memories
- Still has low false positive rate when person denies knowledge

Virtual Reality Mock Crime

- Subjects received email detailing their “Mission”
- Sneak into graduate student office to break in to virtual apartment
- Apprehended and interrogated using ERP-based procedure
- Some subjects given details about utilizing countermeasures
- Innocent subjects tour the same virtual apartment, but with different objects and details.



ARIZONA DAILY WILDCAT Let's smoke pot

POLICE BEAT

Police Beat
By David Halperin
Arizona Daily Wildcat
Friday December 6, 2002

Suspicious e-mail sent

An employee reported that he received an e-mail Wednesday stating he is supposed to commit a crime today, reports stated.

At about 11:35 a.m., the employee told police he had received the suspicious e-mail while in his office at the Arizona Health Sciences Center, 1501 N. Campbell Ave.

The employee told police he did not know the sender of the message or why he received it. He decided to report the incident after his supervisor advised him to do so.

The message read: "This message is simply a reminder of the crime you are to commit on December 6th at 9:00a.m. You should have carefully read over your mission plan and memorized all relevant information in order to carry out your mission. Remember, do not bring materials with you related to the crime and maintain your innocence at all times. Good luck. Dispose of this message once understood," reports stated.

Results of Mock Crime Brainwave Procedure

Group	N	Verdict	
		Guilty	Innocent
Guilty	15	47%	53%
Guilty (countermeasure)	45	17%	83%
Innocent	15	6%	94%

Note: Using Bootstrapping approach, Guilty detection drops to 27%, but innocent subjects classified correctly in 100% of cases. Allows indeterminate outcomes

ERPS and Affective Processing

- IAPS = International Affective Picture System
 - Pleasant, Neutral, Unpleasant
 - Vary in Arousal: Pleasant and Unpleasant tend to be more arousing
- Predict more significant stimuli produce larger P3





Fig. 1. Stimulus synchronized grand average ERP waveforms for Fz, Cz, and Pz electrodes during viewing of affective pictures, separately for each valence category (pleasant, neutral and unpleasant). The left panel illustrates the picture onset potentials on a finer time scale, and the vertical line at Pz illustrate the time area subjected to statistical analysis (i.e. 200–300, 300–400, 400–700, 700–1000 ms). The right panel shows the subsequent 5 s of slow potential change.

Long (6 sec) Presentation Duration


Schupp et al (2000), Psychophysiology



1.5 sec Presentation Duration


Cuthbert et al (2000), Biological Psychology

Figure 1. Picture onset synchronized grand-average event-related potential (ERP) waveforms for each valence category (pleasant, neutral, and unpleasant) from midline electrodes Fz, Cz, and Pz.



120 msec Presentation Duration

Schupp et al (in press), Psychophysiology



ERPS and Implicit Affective Processing

- Ito & Cacioppo (2000) JESP
 - Evaluative Processing (positive vs negative)
 - Nonevaluative (people vs no-people)




FIG. 2. Averaged event-related potential waveforms at electrode Pz as a function of target and context valence. The top panel depicts explicit evaluative categorization effects data from participants in the evaluative task condition. The bottom panel depicts implicit evaluative categorization effects (data from participants in the nonevaluative task condition). The late-positive potential is the positive (downward) deflection peaking at approximately 450–550 ms.

Ito & Cacioppo (2000) *JESP*