


A wee bit more Cardiovascular Psychophysiology ...and then...

The Skeletomotor System

Announcements 2/29/16

- Happy Leap Day!
- Electricity test – See me about retake if needed
- Lab: Get your scored data done by tomorrow!
- 3x5 time

Last session's 3-by-5's (and other inquiries)



Returning to last time

BP and Stress?

Psychophysiology, 47 (2009), 327-332. Blackwell Publishing Inc. Printed in the USA.
Copyright © 2009 Society for Psychophysiological Research
DOI: 10.1111/j.1469-8886.2007.08622.x

Cardiovascular effects in adolescents while they are playing video games: A potential health risk factor?

PETER BORUSIAK,^a ANASTASIOS BOUKIDIS,^a RÜDiger LIERSCH,^a AND JARROD B. RUSSELL^{b,c}

^aZentrum für Kinder- und Jugendmedizin, Helios Klinikum Wuppertal, Wuppertal, Germany

^bZentrum für Kinder- und Jugendmedizin, Evangelisches Krankenhaus Oberhausen, Oberhausen, Germany





Fig 1. Study design (RS1, RS2 = resting phase 1 and 2; P1, P2 = video game phase 1 and 2; arrows indicating blood pressure measuring)


BP and Stress?



Differences were significant for heart rate (HR), systolic (SBP) and diastolic blood pressure (DBP) comparing resting phases and game phases. No significant differences could be found comparing RS1 vs. RS2 and P1 vs. P2 (RS1, RS2 = resting phase 1 and 2; P1, P2 = video game phase 1 and 2).

- Significantly elevate BP during Video Game (VG)
- Energy consumption during Video Game unaltered compared to Rest, and significantly lower compared to Exercise!
- "Comparing all measured parameters it can be said that the relation of blood pressure and energy consumption during VG might not be favorable."

Ballistocardiography





Cardiac Vagal Control and Modulation

- Two Vagal Efferent Branches which terminate on SA Node (Porges 1995, 2003, 2007)
 - Reptilian “Dumb”: Dorsal Motor Nucleus
 - Massive reduction in HR & conservation of oxygen.
 - Dive reflex – cold water on the face during breath hold
 - Phylogenetically newer “smart” Vagus
 - Originates from Nucleus Ambiguus
 - Modulates influence to:
 - Promote attentional engagement, emotional expression, and communication.
 - Mobilizes organism to respond to environmental demands
 - Phasically withdraws inhibitory influence, increasing HR
 - Upon removal of the environmental stressor, resumes its efferent signal
 - Slowing heart rate
 - Allows the organism to self-sooth
- This polyvagal theory is not without its critics (e.g., Grossman & Taylor, 2007).

Measuring Vagal Influence

- Descending Vagal Influence slows HR
- Respiration interrupts this vagal influence
- The size of periodic oscillations due to respiration can therefore index the strength of the Vagal influence
 - Note, however, that under some circumstances, there can be dissociation between RSA and presumed central cardiac vagal efferent activity (cf., Grossman & Taylor, 2007)
- Concerns over changes in rate, and to lesser extent depth
- See special issue of *Biological Psychology*, 2007 for more in depth treatment of these issues and more!



Bradycardia observed in a diving seal. Data adapted from R.S. Elsner (1998), courtesy of <http://www.deeperblue.net/article.php/225>

| | ANS Component | Behavioral Function | |
|-----|--|--|-----------------------------------|
| III | Myelinated vagus (ventral vagal complex) | Social communication, self-soothing and calming, inhibit "arousal" | Nucleus ambiguus |
| II | Sympathetic-adrenal system | Mobilization (active avoidance) | Spinal cord |
| I | Unmyelinated vagus (dorsal vagal complex) | Immobilization (death feigning, passive avoidance) | Dorsal motor nucleus of the vagus |


Fig. 1. Phylogenetic stages of the polyvagal theory.

Porges, 2007

Tonic Vs Phasic


- Tonic Level indexes capacity
- Phasic change indexes actualization of that capacity
- Attention
 - higher vagal "tone" was associated with faster reaction time to a task requiring sustained attention
 - Hyperactive kids treated with Ritalin (Porges, Walter, Korb, & Sprague, 1975).
 - attentional skills improved
 - appropriate task-related suppression of heart rate variability was observed while performing the task requiring sustained attention
- Emotion
 - Beauchaine (2001):
 - low baseline vagal "tone" is related to negative emotional traits
 - high vagal withdrawal is related to negative emotional states

Task-related and Emotion-related modulation



Movius & Allen, 2005

Vagal Control and Defensive Coping



Movius & Allen, 2005

Individual Differences in Cardiac Vagal Control (aka "Trait Vagal Tone")

- Infants
 - Various sick infants have lower vagal tone (Respiratory Distress Syndrome, Hydrocephalic)
 - Infants with higher vagal tone (Porges, various years)
 - More emotionally reactive (both + & -)
 - More responsive to environmental stimuli (behaviorally and physiologically)
- Anxiety Disorders
 - Lower Vagal Tone in GAD (Thayer et al., 1996)
 - Lower Vagal Tone in Panic Disorder (Friedman & Thayer, 1998)
- Depression
 - Depression characterized by lower Vagal tone?
 - State dependent? (Chambers & Allen, 2002)



BIOL PSYCHIATRY
1996;39:255-266 259

Figure 1. Power in the high frequency (respiratory) component of heart period variability in GAD patients and controls during relaxation and worry.

Table 1
Significant contrasts among panickers, blood phobics, and controls

| Variable | Panic (mean, S.D.) | Blood phobic (mean, S.D.) | Control (mean, S.D.) | T ratio, df, p value |
|--|--------------------|---------------------------|----------------------|---|
| IBI (ms) | 761.8 (141.0) | 837.1 (92.4) | 905.2 (132.5) | P < B 4.59 (215) p < 0.001 P < C 7.65 (214) p < 0.001 B < C 4.30 (207) p < 0.001 |
| VAR (ms ²) | 3942 (4009) | 4334 (2663) | 6112 (4563) | P < C 3.70 (214) p < 0.001 B < C 3.44 (207) p < 0.001 P < B N.S. |
| MSD (ms) | 44.4 (31.2) | 55.6 (22.7) | 71.4 (32.1) | P < B 3.05 (215) p < 0.001 P < C 6.34 (214) p < 0.001 B < C 4.11 (207) p < 0.001 |
| HF power (ms ² Hz ⁻¹) | 991 (1225) | 1385 (1073) | 2239 (1911) | P < C 2.49 (212) p < 0.05 P < C 5.67 (212) p < 0.001 B < C 3.90 (203) p < 0.001 |
| LF/HF | 2.1(2.5) | 1.3 (1.8) | 1.0 (1.5) | P < C 2.41 (209) p < 0.05 P < C 3.64 (203) p < 0.001 B < C N.S. |

P, panic; B, blood phobic; C, control.

Chambers and Allen (2002) *Psychophysiology*

Can Vagal Control predict development of anxiety following stressors?




Fig. 1. Effect of the interaction between RSA adjusted for age and Time since initial assessment on TMAS over a 1-year period. Although RSA is a continuous variable, for illustrative purposes, its effect on TMAS is plotted at ± 1 SD from the mean. Error bars represent standard errors. RSA: respiratory sinus arrhythmia; SD: standard deviation; TMAS: Taylor Manifest Anxiety Scale.

Kogan, Allen, Wehhs (2012)
Biological Psychology

Trait Vagal Tone as Moderator of Response following Bereavement

- Bereavement as a period of cardiovascular risk
- Disclosure as an intervention for Bereavement (O'Connor, Allen, Kaszniak, 2005)
- Overall, all folks get better, but no differential impact of intervention
- BUT... Vagal Tone as moderator





Figure 2. Scatterplot, prediction line, and prediction equation for the relationship between respiratory sinus arrhythmia (log of the variance of the band-limited (12–40 Hz) HRV series) and physical health complaint (negative physical health complaints) for the disclosure group (top panel) and the control group (bottom panel). Negative physical health complaint score represents improvement from baseline to follow-up.

Orienting, Attention, and Defense *Emotional reactivity*






Figure 3. Scatterplot, prediction line, and prediction equation for the relationship between respiratory sinus arrhythmia (log of the variance of the band-limited (12–40 Hz) HRV series) and physical health complaint (negative physical health complaints) for the disclosure group (top panel) and the control group (bottom panel). Negative physical health complaint score represents improvement from baseline to follow-up.



OR Vs DR

Electromyography

Why Record EMG?

- Facial Musculature rich; emotional expressions; a “leaky channel of expression”
- Startle blink as a probe for affective valence
- Muscle tension in disorders and stress
- Record “pre-behavioral” motor output
 - Facial Expressions
- Human Performance (e.g incorrect channel EMG in forced-choice RT task)

The Expressive Face

- [Clip 1](#)
- [Clip 2](#)

Striated Muscle

- Large number of muscle fibers arranged in parallel
- “Striated” reflects that these fibers actually comprise smaller fibrils
 - Fibrils have repeating cross striations (Z-lines)
 - Fibrils plus tissue between = Sarcomeres

Striated Muscle




Figure 12.2. Diagram of the structure of the muscle with increasing magnification going from left to right. The bottom right corner of the figure illustrates the microgeometric changes that occur with contraction. (Modified from Figure 10.7 of Schmidt-Nielsen, 1997).

Striated Muscle

➤ During contraction:

- Very small changes in length of filaments
- But big changes in the distance between the Z-bands as the thick filaments slide between the thin

Innervation

- Muscle needs stimulation to contract
- The motor nerve
 - Contains many motoneurons
 - Each motoneuron branches into several axon fibrils
- At end of each axon fibril is a junction with the muscle fiber
 - Known as the motor endplate




Figure 12.3. Diagram of two motor units. (Modified from slide 3705 of Netter, 1991).

Innervation

- Each motoneuron innervates several to many muscles (innervation ratios 10:1 to 2000:1), but each muscle innervated by only one motoneuron
- Therefore, muscle fibers fire simultaneously or in concert with one another
- Stronger contractions due to either more motoneurons firing, or increases in rate of already firing motoneurons





Figure 12.3. Diagram of two motor units. (Modified from slide 3705 of Netter, 1991).

Cartoon of how it works



Before






After









What is EMG signal?

- Reflects electrical field generated by Muscle Action Potentials (MAPs)
- Small portion conveyed to surface via extracellular fluids to skin
- Can also record invasively with subcutaneous needle electrodes





EMG Power



Signal Recording

- MAPs summate in quasi-random fashion to produce resultant signal
 - Range of ~10-500 Hz
 - Amplitude of sub-microvolt to over 1000 microvolts
- Note overlap with 60 Hz range
 - Prepare ground site carefully; Differential amplifier will assist in removing 60 Hz
 - Prepare recording sites carefully to lower impedance
 - Shielded rooms and leads can help
 - Can also filter out this range, but may toss “baby with bathwater”

Signal Recording (cont')

- Can use wide variety of electrodes
 - Ag-AgCl still preferred
 - Small size increases specificity of recording
- Skin Prep
 - Abrade to reduce impedance to $< 5\text{ K}\Omega$
- Use Bipolar arrangements, in line with long direction of muscle of interest
- Use common ground for all sites
- Keep wires and such out of subject’s visual field
- Describe placements precisely
 - Standard for location is Fridlund & Cacioppo (1986) for facial EMG placements

Signal Recording (cont')

- Amplification
 - Differential amplifiers with common mode rejection
 - Actually double differential (ground)
- Amplify voltages 1000-20000 times
- May use on-line filter
 - Should pass 10-500 Hz
- Digitization (more in next lecture)
 - Fast, very fast
 - Or, slower, following on-line signal processing






Figure 12.5. Common alternative representations of the surface EMG signal. The top five smaller panels depict three distinct non-linearities: raw (1), band-pass filtered only (2), before the second half-wave rectified waveform, the third, full-wave rectified waveform, and the fourth, integrated waveform. The larger bottom panel depicts what one does if one wants to look at the signal in the frequency domain. (Modified from Figure 7 of Cacioppo et al., 1993.)


Corrugator “Frown”



Zygomatic “Smile”




Looking at PICTURES




A few Applications

- Startle Probe
- Subtle affect
 - Mere Exposure
 - Subliminal effects
 - Mortality Salience
 - Biofeedback of EEG -- outcome measure
 - Emotion Regulation – outcome measure
 - Empathy – individual difference measure




► Neural Circuits Responsible for an Auditory Startle Response and for Its Augmentation by Conditioned Aversive Stimuli





Source: Adapted from Davis, M., Trends in Pharmacological Sciences, 1992, 13, 35–41.


International Affective Picture System (IAPS)




Affective Space:
Picture Content and Gender



Corrugator EMG



Zygomatic EMG



Resting HRV as moderator of Startle Potentiation




Figure 1. Mean startle amplitude as a function of baseline HRV and valence. Startle amplitudes are in microvolts.

From: Ruiz-Padiala, Sollers, Vila, & Thayer (2003) *Psychophysiology*

A few Applications

- Startle Probe
- Subtle affect
 - Mere Exposure
 - Subliminal effects
 - Mortality Salience
 - Biofeedback of EEG -- outcome measure
 - Emotion Regulation – outcome measure
 - Empathy – individual difference measure

The Phenomenon:

- People prefer stimuli to which they have been previously exposed to unfamiliar stimuli
- In absence of any reinforcement (“mere” exposure)
- Examples:
 - People we see incidentally in our routines
 - Songs
 - Scientific journal preferences
- Effect size $r=.26$ (Meta-analysis, Bornstein, 1989)

The logic:


- Evolutionary account Bornstein (1989)
 - it may be adaptive to prefer the familiar over the novel
 - novel objects could present a potential threat
 - organisms that had a fear of the strange and unfamiliar were more likely to survive, reproduce, and pass on genetic material
 - Preferring the familiar may thus be an adaptive trait that has evolved in humans and nonhumans
- Prediction:
 - unfamiliar as compared with familiar stimuli may be associated with more negative attitudes because of the unfamiliar stimuli's association with potential danger
 - Thus may see greater corrugator activity to novel than to familiar
 - No prediction for positive affect (Zygomaticus activity)



A few Applications

- Startle Probe
- Subtle affect
 - Mere Exposure
 - Subliminal effects
 - Mortality Salience
 - Biofeedback of EEG -- outcome measure
 - Emotion Regulation – outcome measure
 - Empathy – individual difference measure

Loosely translated from Harmon-Jones & Allen, 2001

Dimberg et al *Psychological Science* 2000

PSYCHOLOGICAL SCIENCE

Unconscious Facial Reactions




Fig. 1. Mean facial electromyographic response for the *zygomatic major* muscle, plotted in intervals of 100 ms during the first second of exposure. Three different groups of participants were exposed to identical neutral faces ("Ne"), preceded by unconscious exposure of happy, neutral ("neutr"), or angry target faces, respectively.





Fig. 2. Mean facial electromyographic response for the *corrugator supercilii* muscle, plotted in intervals of 100 ms during the first second of exposure. Three different groups of participants were exposed to identical neutral faces ("Ne"), preceded by unconscious exposure of angry, neutral ("neutr"), or happy target faces, respectively.

Dimberg et al *Psychological Science* 2000


A few Applications

- Startle Probe
- Subtle affect
- Mere Exposure
- Subliminal effects
- Mortality Salience
- Biofeedback of EEG -- outcome measure
- Emotion Regulation – outcome measure
- Empathy – individual difference measure



A few Applications

- Startle Probe
- Subtle affect
- Mere Exposure
- Subliminal effects
- Mortality Salience
- Biofeedback of EEG -- outcome measure
- Emotion Regulation – outcome measure
- Empathy – individual difference measure



From Allen, Harmon-Jones, and Cavender (2001)
Allen, Cavender, Harmon-Jones, *Psychophysiology* 2001

A few Applications

- Startle Probe
- Subtle affect
 - Mere Exposure
 - Subliminal effects
 - Mortality Salience
 - Biofeedback of EEG -- outcome measure
 - Emotion Regulation – outcome measure
 - Empathy – individual difference measure




Figure 1. Self-reported negative affect on a 7-point Likert scale, where 0 = "not negative at all" and "7" = "strongly negative."

Ray, McRae, Ochsner, & Gross, *Emotion*, 2010




Figure 2. Standardized (A) corrugator EMG and (B) startle magnitude (averaged over Times 1 and 2).
Ray, McRae, Ochsner, & Gross, *Emotion*, 2010

A few Applications

- Startle Probe
- Subtle affect
 - Mere Exposure
 - Subliminal effects
 - Mortality Salience
 - Biofeedback of EEG -- outcome measure
 - Emotion Regulation – outcome measure
 - Empathy – individual difference measure




Figure 1. The zygomatic major muscle response to pictures of happy and angry facial expressions for the High and Low empathy groups, plotted as a function of 100-ms intervals during the first second after stimulus onset.




Figure 2. The corrugator supercilii muscle response to pictures of happy and angry facial expressions for the High and Low empathy groups, plotted as a function of 100-ms intervals during the first second after stimulus onset.
Dimberg & Thunberg (2012) *PsyCh Journal*