



## The Event-Related Potential (aka the ERP) (Part 2)



## Applications of Early Components

- Neurological evaluation of sensory function; e.g. evaluation of hearing in infants
- Tones of various dB intensities presented and V wave in auditory brainstem ERP examined
- **Figure 10**; 4000 individual trials per average

## Prediction of recovery from coma




- ❑ Somatosensory evoked potentials were recorded from a patient who was still comatose 1 week after severe closed head injury.
- ❑ Responses evoked by electrical stimulation of left and right median nerves
- ❑ Normal tracing seen at Erb's point, and from the next over vertebra prominens, but not over C3' of C4'.
- ❑ Absence of any cortical response a bad prognostic sign. Patient continued in a chronic vegetative state 1 year after accident

## Inter-Hemispheric Transfer Time (IHTT)

- Hypothesized that interhemispheric transfer of information may be abnormal in various disorders (e.g., dyslexia)
- Reaction Time measures contain too much variability not related to Transfer Time
- ERP early components appear promising as a measure of time required to transfer information between hemispheres

## IHTT Study

- Checkerboards subtending < 1 degree of visual angle presented 2.9 degrees from center
- ERP's recorded at O1 and O2
- Problem of lateralization and Paradoxical results possible; parafoveal regions on banks of calcarine fissure
- P100 wave latency examined; earlier latency in occiput contralateral to presentation
  - Measured by peak picking procedure
  - Also by cross-lagged correlation technique
  - Both methods suggest ~15 millisecond IHTT; found to be in expected direction predicted by anatomy for over 90% of subjects
  - Reaction time data from same task showed no reliable differences



Saron & Davidson, 1989





Fig. 1. Paradigm for using ERPs to study attention. Stimulus display (left) and idealized results (right). Subjects fixate a central cross and attend either to the left or right visual field. Stimuli are then presented to the left and right visual fields in a rapid sequence. In this example, the ERP elicited by a left visual field stimulus contains larger P1 and N1 components when the stimulus is attended ('Attend left') than when it is ignored ('Attend right').

From Luck et al, *TICS*, 2000

## More than Spatial Directed Attention




Fig. 2. Grand averaged visual ERPs at Pz electrode for the 3 array sizes, showing the shorter latencies, larger P1s for array size 17, but longer latency P3 (dark arrows) than for array sizes 5 and 9 (grey arrows). These are averaged across colour, orientation and conjunction conditions, as this ERP effect was seen regardless of whether it was a single feature or conjunction trial.

Taylor  
*Clinical Neuropsych*  
2002

## More than Spatial Directed Attention





Fig. 3. Mean P1 latencies across 7 age groups, showing the consistently shorter latencies to faces compared to inverted faces and control stimuli (phase-scrambled faces and flowers). There were 15 children in each of the 6 age groups and 38 adults (adapted from Taylor et al., 2001c).

Taylor  
*Clinical Neuropsych*  
2002

"These combined PET/ERP data therefore provide strong evidence that sustained visual spatial attention results in a preset, top-down biasing of the early sensory input channels in a retinotopically organized way"



Woldorff et al., *Human Brain Mapping*, 1997

## Prelude to Advance Topic: Source Localization




Figure 3. Left: Observed potential distributions in the attend-left-minus-attend-right difference waves at the peak of the P1 attention effect (110–130 msec). Right: Corresponding model potential distributions seeded by the dorsal occipital PET foci which provided an excellent fit to the P1 effect (residual variance 2%).



## Information Transmission







Fig. 2. Grand averaged visual ERPs at Pz electrode for the 3 array sizes, showing the shorter latencies, larger P1s for array size 17, but longer latency P3 (dark arrows) than for array sizes 5 and 9 (grey arrows). These are averaged across colour, orientation and conjunction conditions, as this ERP effect was seen regardless of whether it was a single feature or conjunction trial.

Taylor  
Clinical Neurophys  
2002

## P3 Latency

- An index of processing time, independent of response requirements
  - RT measures confounds the two
  - McCarthy & Donchin (1981) experiment:
    - The words "RIGHT" or "LEFT" embedded in a matrix of letters of X's
    - Compatible condition: respond with hand indicated in matrix; Incompatible condition: respond with opposite hand (e.g., LEFT signals right hand response);
    - Results:
      - P300 latency delayed when discriminability more difficult
      - Response compatibility had no effect on P300 latency
      - **Note amplitude reduction as function of noise-information transmission)**



Not only difficulty in physical discrimination, but difficulty in cognitive categorization


Figure 4.10. ERP waveforms at Pz averaged across subjects for three different semantic categorization tasks. The solid line indicates ERPs obtained during a task in which the subjects had to distinguish between the word DAVID and the word NANCY (the FN condition). The dotted line indicates ERPs obtained during a task in which the subjects had to decide whether a word presented was a male or a female name (the VN condition). The dashed line indicates ERPs obtained during a task in which the subjects had to decide whether a word was or was not a synonym of the word PROD (the SYN condition). These three tasks were considered to involve progressively more difficult discriminations. Note the latency of P300 peak is progressively longer as the discrimination is made more difficult. (Copyright 1977, AAAS. Adapted with permission of the author and publisher from Kutas, McCarthy, & Donchin, 1977.)

## Construct Validity?

- What, then, does the P300 mean in very general terms?
  - A stimulus (or class of stimuli) is "important"; denotes information that is necessary or useful to the task
  - Stimulus is meaningful, important, noticeable
  - Evaluated within context of working memory? (cf. Donchin & Coles, 1988; Verleger 1988; Polich, 2007; Verleger, 2008)
- The P3a (Squires, Squires, and Hillyard, 1975): P3-like component with a frontal maximum and occurs to improbable stimuli in the "to-be-ignored" class of stimuli; a novelty response.

## How Many P3s?

- The Classic P3/P300
  - Parietal Central Maximum
  - Largest when stimuli rare and task-relevant
- The P3a (Squires et al., 1975) or Novelty P3 (Courchesne et al., 1975)
  - More anterior scalp distribution
  - Slightly earlier latency
  - Responsive to rare, unexpected, unattended stimuli
    - Courchesne: "deviant non-target stimuli: buzzes, filtered noises and other unusual sounds"



Simons et. al, 2001

- Squires Task was tones (two tones)
- Courchesne task was digitized speech ("me" "you" and collection of naturally occurring sounds)
- In all cases subjects merely counted Tones

Fig. 1. ERP waveforms (left) and PCA basis waves (right) obtained from infrequent targets during the Squires (top) task and infrequent nontargets/nerths during the Courchesne (bottom) task. PCA was conducted during the 220–420 ms epoch following stimulus onset and four factors were extracted from each data set.


## P3a – Can you see it?

- Some inconsistencies in finding P3a following the initial Squires, Squires and Hillyard 1975 report
- Comerchero & Polich (1998) may have resolved the enigma
  - P3a highly dependent on foreground discrimination



Table 1  
Stimulus type (probability) for each task condition and modality (auditory = frequency and intensity, visual = area and shape-color)

Modality	Auditory		Visual	
Nontarget distinctiveness	Low	High	Low	High
Target (0.10)	2000 Hz	2000 Hz	12.57 cm <sup>2</sup>	12.57 cm <sup>2</sup>
	75 dB	75 dB	● Blue	● Blue
Standard (0.80)	1940 Hz	1940 Hz	10.18 cm <sup>2</sup>	10.18 cm <sup>2</sup>
	75 dB	75 dB	● Blue	● Blue
Nontarget (0.10)	500 Hz	4000 Hz	12.57 cm <sup>2</sup>	12.57 cm <sup>2</sup>
	75 dB	90 dB	■ Blue	■ Fuchsia



Note: Nontarget peak amplitude was earlier and larger at the frontal electrodes than those from the target stimuli, but especially when foreground discrimination is difficult

Comerchero & Polich (1998).  
*Clinical Neurophysiology*




Fig. 1. Schematic illustration of the single-stimulus (top), oddball (middle), and three-stimulus (bottom) paradigms, with the elicited ERPs from the stimuli of each task at the right (Polich and Criado, 2006). The single-stimulus task presents an infrequent target (T) in the absence of any other stimuli. The oddball task presents two different stimuli in a random sequence, with one occurring less frequently than the other target = T, standard = S). The three-stimulus task is similar to the oddball with a compelling distractor (D) stimulus that occurs infrequently. In each task, the subject is instructed to respond only to the target and otherwise to refrain from responding. The distractor elicits a P3a, and target elicits a P3b (P300). Reprinted with permission of the authors and from Elsevier (Copyright 2006).

Polich, *Clin Neurophys*, 2007

## Synopsis

“...the manipulation of target-standard stimulus discriminability produced a stimulus environment in which the infrequently occurring nontarget engaged focal attention in a manner similar to that observed previously for ‘novel’ stimuli.”

“However, all stimuli in the present study were employed because of their ‘typical’ characteristics, so that the results imply that an anterior P3a component can be produced without using ‘novel’ stimuli per se.”

“If stimulus context is defined primarily by a difficult target/standard discrimination, attentional redirection to the nontarget would occur because of the frontal lobe activation that generates P3a.”

Comerchero & Polich 1998, p. 47

## ERPs and Memory

- Sensitive to both Recognition
  - Likely episodic recollection
- Sensitive to Encoding

## Repetition Priming Effects

- Robust effect that repeated items produce an enhanced late positivity across a broad latency range
- Magnitude of effect related to strength of memory trace






Fig. 4. Grand mean ERP waveforms elicited by correctly recognized old and correctly rejected new items from Johnson et al. (1998a). The left column depicts the old and new waveforms at the electrode site and hemisphere where that subcomponent was largest. Reproduced from Johnson et al. (1998a) with permission of the publisher.

## Repetition Priming

- Are there repetition effects that do not depend on the subjective awareness of the subject?
  - Can use Masked Priming to examine (Schnyer, Allen, Forster, 1997)




Schnyer, Allen, Forster, 1997




Standard Repetition Effect for Words Seen Unmasked in Previous Blocks  
Task is to make OLD-NEW decision

Schnyer, Allen, Forster, 1997



Standard Repetition Effect for Words Seen Unmasked in Previous Blocks  
But Task is to make WORD-NONWORD decision

Schnyer, Allen, Forster, 1997




Masked Repetition Priming Effect for Words Presented only a Trial Previously

Schnyer, Allen, Forster, 1997

## Memory Encoding

- Words subsequently remembered show enhanced positivity at encoding
- Strategy interacts, however



Note prototypic DM effect on left, but not on right for those that used elaborative strategies. Note enhancement over frontal lead for these latter subjects.

Figure 4.12. ERPs elicited by "isolated" words that were later recalled (solid line) or not-recalled (dashed line). The left column shows ERPs for subjects who used rote mnemonic strategies; the right column shows ERPs for subjects who used elaborative strategies. Note that the amplitude of P300 is related to subsequent recall for the rote memorizers, but not for elaborators. (Copyright 1986, Elsevier Science Publishers. Reprinted with permission of the publisher from Fabiani, Karis, & Donchin, 1986b.)




Fig. 3. A: Grand mean ERPs elicited by study items that were subsequently associated with remember or know judgments (Modified from Friedman and Trott, 2000). C: CSD maps for 2 intervals (500-800; 810-1100 ms) measured in the Dm waveform associated with a subsequent Remember judgment. Data in A and B recorded at a left inferior prefrontal scalp site.

## Indirect Assessments of Recognition

- Can the ERP detect recognition, independent of subjects' overt responses?
- Two applications
  - Clinical Malingering
  - Forensic Assessment


## ERP Memory Assessment Procedures

- Learn a list of words
- Learn a second list of words
- Task: Concealed (1<sup>st</sup> list) and Nonconcealed (2<sup>nd</sup> list) words appear infrequently

Item Type	Probability	Response	P3 Amplitude
Nonconcealed	1/7	"Yes"	Large
Concealed	1/7	"No"	Large if Recognized Small if not Recognized
Unlearned	5/7	"No"	Small


- Similar to procedures by Rosenfeld et al, Farwell & Donchin

## The Classic Oddball Experiment



## Motivational Variations

Conceal	Lie	Lie + \$\$
<ul style="list-style-type: none"> <li>➤ "YES" for words <u>JUST</u> learned, "NO" for all others</li> <li>➤ Try to hide the fact that you learned the first list of words I taught you</li> </ul>	<ul style="list-style-type: none"> <li>➤ "YES" for words learned</li> <li>➤ Lie about words from the first list I taught you</li> </ul>	<ul style="list-style-type: none"> <li>➤ "YES" for words learned</li> <li>➤ Lie about words from the first list I taught you</li> <li>➤ \$5.00 incentive</li> </ul>




After Allen & Iacono, 1997

## The Challenge

*To provide statistically supported decisions for each and every subject, despite considerable individual variability in ERP morphology*





### Bayesian Combination of ERP Indicators: Probability that an ERP was elicited by Learned Items

Subject	List						
	Learned		Unlearned				
	NonConceal	Conceal	U1	U2	U3	U4	U5
#01	1.0	0.999	0.000	0.000	0.000	0.000	0.001
#02	1.0	1.0	0.000	0.000	0.000	0.000	0.000
#03	1.0	0.999	0.000	0.000	0.000	0.002	0.000
#04	1.0	1.0	0.000	0.001	0.002	0.000	0.000
#05	1.0	0.971	0.002	0.000	0.000	0.000	0.000
#06	1.0	0.999	0.000	0.000	0.000	0.000	0.000
#07	0.983	1.0	0.000	0.000	0.000	0.000	0.000
...							
#18	0.996	0.983	0.874	0.001	0.000	0.000	0.000
#19	0.009	0.214	0.971	0.000	0.002	0.189	0.983
#20	1.0	0.999	0.002	0.000	0.009	0.000	0.214

Note: Only trials in which subjects did not acknowledge concealed items included

### Classification Accuracy based on ERPs

	Learned (true pos)	Unlearned (true neg)
Conceal	0.95	0.96
Lie	0.93	0.94
Lie + \$\$	0.95	0.98
Combined	0.94	0.96


Allen, Iacono, & Danielson, *Psychophysiology*, 1992

### Extensions from Lab to Life...


- Two tests of the robustness of this procedure:
  - False recollections
  - Virtual Reality Mock Crime

### A Laboratory Paradigm for False Recollections: DRM


- Subjects presented with 15 words highly associated with an omitted critical item



### Reported Rates of Recognition



Allen and Mertens (2008)




## The Box Score Blues

Ground Truth	Test Verdict
	Recognized
Actually Learned	56%
Critical Lure	72%
Unlearned	4%

- ❑ Highlights the need to have memorable items in the test
- ❑ Suggests limited utility in substantiating disputed memories; e.g., claims regarding recovered memories
- ❑ Still has low false positive rate when person denies knowledge

## Virtual Reality Mock Crime

- Subjects received email detailing their “Mission”
- Sneak into graduate student office to break in to virtual apartment
- Apprehended and interrogated using ERP-based procedure
- Some subjects given details about utilizing countermeasures
- Innocent subjects tour the same virtual apartment, but with different objects and details.




## Results of Mock Crime Brainwave Procedure

Group	N	Verdict	
		Guilty	Innocent
Guilty	15	47%	53%
Guilty (countermeasure)	45	17%	83%
Innocent	15	6%	94%

Note: Using Bootstrapping approach, Guilty detection drops to 27%, but innocent subjects classified correctly in 100% of cases. Allows indeterminate outcomes


## ERPs and Affective Processing

- IAPS = International Affective Picture System
  - Pleasant, Neutral, Unpleasant
  - Vary in Arousal: Pleasant and Unpleasant tend to be more arousing
- Predict more significant stimuli produce larger P3



Long (6 sec)  
Presentation Duration

Schupp et al (2000),  
*Psychophysiology*




1.5 sec Presentation  
Duration

Cuthbert et al (2000),  
*Biological Psychology*

Fig. 1. Stimulus synchronized grand average ERP waveforms for Fz, Cz, and Pz electrodes during viewing of affective pictures, separately for each valence category (pleasant, neutral and unpleasant). The left panel illustrates the picture onset potentials on a finer time scale, and the vertical lines at Pz illustrate the time areas subjected to statistical analysis (i.e. 200–300, 300–400, 400–700, 700–1000 ms). The right panel shows the subsequent 5 s of slow potential change.

Figure 1. Picture onset synchronized grand-average event-related potential (ERP) waveforms for each valence category (pleasant, neutral and unpleasant) from midline distribution Fz, Cz, and Pz.



120 msec Presentation  
Duration

Schupp et al (2004),  
*Psychophysiology*




Figure 8. Sensor outline of the generic sensor net. The left and right panels illustrate the sensor clusters used to quantify the early (ERP) and late (LPP) relative ERP components, respectively.

## ERPs and Implicit Affective Processing

### ➤ Ito & Cacioppo (2000) *JESP*

- Evaluative Processing (positive vs negative)
- Nonevaluative (people vs no-people)





FIG. 2. Averaged event-related potential waveforms at electrode Pz as a function of target and context valence. The top panel depicts explicit evaluative categorization effects data from participants in the evaluative task conditions. The bottom panel depicts implicit evaluative categorization effects data from participants in the nonevaluative task condition. The late-positive potential is the positive (downward) deflection peaking at approximately 450–550 ms.


Ito & Cacioppo (2000) *JESP*

## N400 and Language



- Originally reported by Kutas & Hillyard, 1980.
- Semantic Incongruity is separable from other forms of deviations (e.g. large font)
  - N400 Semantic Deviation
  - P300 Physical Deviation
- Also seen in semantic differentiation tasks (Polich, 1985); APPLE, BANANA, ORANGE, MANGO, TRUCK
- Subject-Object mismatch (the Florida group)
- NOTE: N400 will appear before P3 (which will be ~P550 in word tasks)

## N400 and Language



Sensitive to degree of semantic incongruity

## Political Evaluations!

➤ Morris Squires et al. *Political Psychology* 2003




Figure 2. Attitude-priming paradigm and examples of its use.

ERPs and Hot Cognition

739

CONGRUENT  
INCONGRUENT





Figure 4. ERPs to congruent and incongruent prime/target pairs.


Congruent or incongruent defined based on idiographic data from pretest

Morris Squires et al. *Political Psychology* 2003




- Cloze probability: proportion of respondents supplying the word as continuation given preceding context
- N400 reflects unexpected word given the preceding context
- This is independent of degree of contextual constraint
- Larger N400
  - Low cloze, Contextual constraint high:
    - The bill was due at the end of the hour
  - Low cloze, Contextual constraint low:
    - He was soothed by the gentle wind
- Smaller N400
  - The bill was due at the end of the month

Kutas & Federmeier, 2011




- Sentence completion
  - Best (expected) ending *small*
  - Unexpected but related *larger*
  - Unexpected and unrelated *largest*
- Categorical relations ... sentence final word is:
  - an expected category exemplar
  - an unexpected, implausible exemplar from the same category as the expected one (related anomalous)
  - from a different category (unrelated anomalous)
- Note multiple modalities of effect, and graded effect in RVF (LH)

Kutas & Federmeier, 2011




- Word Association, with second word in pair
  - Unrelated to first (*eat door*)
  - Weakly related to first (*eat spoon*)
  - Strongly related to first (*eat drink*)
- Orthographic neighborhood size (among a list of words, pseudowords, and acronyms)
  - Words that share all but one letter in common with particular word
  - Large 'hood (e.g., *slop*) – *large* N400
  - Small 'hood (e.g., *draw*) – *small* N400

Kutas & Federmeier, 2011



- Math: (e.g.,  $5 \times 8 = \underline{\quad}$ )
  - Correct (40) *small*
  - Related (32, 24, 16) *small if close*
  - Unrelated (34, 26, 18) *large*
- Movement and Gestures
  - Typical actions (cutting bread with knife) = *small*
  - Purposeless, inappropriate, or impossible actions = *large*
    - Cutting jewelry on plate with fork and knife
    - Cutting bread with saw
- N400 modulated by both:
  - appropriateness of object (e.g., screwdriver instead of key into keyhole)
  - features of motor act per se (e.g., orientation of object to keyhole)


Kutas &amp; Federmeier, 2011



- Repetition effects
  - Repetition creates contextual familiarity, reduced processing demands
  - N400 thus useful in studying memory
  - Appears additive with incongruity effects

Kutas &amp; Federmeier, 2011


## N400 – The Unexpected Hero!



## Response-locked potentials

- Lateralized Readiness Potential (LRP), a special case of movement-related potentials
- Error-related Negativity (ERN, aka  $N_E$ )


148 EIMER Eimer 1998, Beh Res Methods



## Lateralized Readiness Potential

- LRP can be stimulus-locked or response-locked
- For stim-locked, latency is time between stimulus onset and LRP onset
- For rps-locked latency is time between an LRP deflection and the overt response.

Figure 1. Computation of the lateralized readiness potential (LRP) with the double subtraction method on the basis of event-related brain potentials (ERP) waveforms elicited at electrodes C3' (left hemisphere) and C4' (right hemisphere). Top panels: Grand-averaged ERP waveforms from 10 subjects elicited at C3' (solid lines) and C4' (dashed lines) in response to stimuli requiring a left-hand response (left side) and to stimuli requiring a right-hand response (right side). Middle panels: Inference waveforms resulting from subtracting the ERPs obtained at C4' from the ERPs obtained at C3' separately for left-hand responses (solid line) and right-hand responses (dashed line). Bottom panel: LRP waveforms resulting from subtracting the C3' - C4' difference waveform for left-hand responses from the C3' - C4' difference waveform for right-hand responses. A downward-going (positive) deflection indicates an activation of the correct response; an upward-going (negative) deflection indicates an activation




## Response conflict in the LRP

Figure 2. Top: Examples of stimulus displays in an experiment or spatial stimulus-response compatibility (Eimer, 1993, Experiment 1a) in which stimulus and response sides could either be compatible (left side) or incompatible (right side). Bottom: Grand-averaged LRP waveforms from 10 subjects, elicited in compatible trials (solid line) and in incompatible trials (dashed line).

Eimer 1998, Beh Res Methods

## The ERN



Flankers Task:

MMNMM

— Error  
----- Correct

Also sometimes termed Ne




Fig. 3. Relationship between error-related negativity (ERN) amplitude and three measures of compensatory behavior. Left panel: Average event-related potentials at the Cz electrode as a function of the four levels of the posterior probability measure of ERN amplitude. Right panel, top: Error squeeze force in Kg as a function of the four ERN levels. Right panel, middle: Probability of error correction as a function of the four ERN levels. Right panel, bottom: Correct reaction time on the trial following an error as a function of the four ERN levels.

Gehring et al., 1993

## Modality Specific?





Fig. 1. Grand averages (Experiment 4,  $n = 12$ ) of the RIA for error (heavy lines) and correct trials (light lines) after visual (vis) and auditory letter stimuli (aud) in a 2x2 task. The error negativity (Ne) is seen as a sharp negative deflection with central maximum peaking at about 100 ms after the incorrect key press (R). The error positivity (Pe) is seen as a late positive deflection with Cz maximum peaking at about 300 ms after the incorrect key press. The correct trials show a positive complex with Fz maximum in seen.

➤ Does not matter what modality stimulus was presented



➤ Does not matter what modality response was made  
➤ Eye

Nieuwenhuis et al., 2001: Saccade Task

C.B. Holroyd et al. / Neuroscience Letters 242 (1998) 65–68




Fig. 2. Source localization of the error-related negativity. Circles represent locations of sources determined for hand and foot responses: (a) coronal view; (b) sagittal view; (c) for comparison, source locations of the ERN determined in previous studies are depicted along with the locations of the ERN obtained in the present study. Squares represent locations of sources found for ERNs elicited by visual, auditory, and somatosensory feedback [10]. Crossed symbols represent locations of sources found for ERNs elicited by errors in two reaction time experiments [2].

➤ Does not matter what modality response was made  
➤ Eye  
➤ Hand  
➤ Foot

## Error Detection Vs. Error Compensation

➤ If Error Compensation, ERN/Ne should not be present in tasks where compensation impossible

➤ Ergo...

➤ the Go-Nogo!

➤ Play along... press only for X following X




Fig. 5. Grand averages (Experiment 2;  $n = 10$ ) of the RTA for false alarms and hits in Go/NoGo tasks (heavy lines), and choice errors and correct choice trials in two-way choice tasks (thin lines). Errors continuous lines, correct responses broken lines. The Ne is delayed relative to the incorrect key press, and the Pe is smaller, for choice errors compared to false alarms. In correct trials a positive complex with Pr maximum is seen, which is larger after visual than after auditory stimuli. However, this complex is not larger for hits than for correct choice trials.

Falkenstein Hoormann Christ & Hohnsbein, *Biological Psychology*, 2000, Summary of Falkenstein et al 1996

## Error Detection Vs. Outcome Impact

- Might the “cost” or “importance” or “salience” of an error be relevant to this process?
- Studies relevant to error salience
  - Speed-accuracy trade off
  - Individual differences

## Speed Vs. Accuracy

M. Falkenstein et al. / *Biological Psychology* 51 (2000) 87–107





Fig. 4. Grand averages (Experiment 1;  $n = 9$ ) of the RTA for correct responses (C), errors (E), and difference wave (E-C) in a 2x2 task under moderate (right lines) and severe time pressure (left lines). The error rates were 15% (moderate) and 30% (severe); the number of error trials used was equalized for the two conditions. The Ne is smaller for severe time pressure/high error rate.

## Individual Differences

- Psychopathy (or analog)
- OCD

## Deficits in Error Monitoring in Psychopathy

- Psychopaths appear unable to learn from the consequences of their errors
  - Avoidance learning deficits
  - In the context of rewards *and* punishments
  - Deficient anticipatory anxiety




Thirty participants selected: 15 high SO

15 low SO


Dikman & Allen, 2000, *Psychophysiology*

## Procedure

- Eriksen flanker task: SSHSS
- Two conditions for each subject
  - Reward (REW), errors “No \$”
  - Punishment (PUN), errors 95 dB tone
- Consequences of errors could be avoided by self-correcting response within 1700 msec window
- Response mapping switched at start of each of 10 blocks, total trials 600
- Only corrected error trials examined



Dikman & Allen, 2000, *Psychophysiology*



## ERN in OCD




Fig. 1. Response-locked event-related potential waveforms at the Cz electrode location. The left panel compares correct-trial and error-trial waveforms for control participants and for individuals with obsessive-compulsive disorder (OCD). The right panel compares error-trial waveforms for the two groups. Times are plotted relative to the latency of the button-press response. ERN = error-related negativity.

And amplitude of ERN correlates with Symptom severity (correlation magnitude  $\sim .50$ ); Gehring et al. (2000)

## Errors and Feedback


- Endogenous Error Detection
- Exogenous Error Feedback
- Common Mechanism?

## Choices and Feedback






## The Feedback Medial Frontal Negativity




Miltner, Braun, & Coles, (1997) *Journal of Cognitive Neuroscience*

## The Gambling Task




Gehring and Willoughby, 2002 *Science*



**Fig. 2.** ERP waveforms, scalp topography, and likely neural generator of the MFN. (A) The waveforms are shown at the Fz (frontal) electrode site. The solid red line corresponds to the average ERP waveform for all trials in which the participant lost money. The dashed green line corresponds to those trials in which the participant gained money. The MFN is indicated by the arrow. The error bar represents two standard errors of the mean, based on the mean squared error from the ANOVA (9). (B) The map of scalp activity shows the voltages, derived by subtracting the loss-trial waveform from the gain-trial waveform, computed at 265 ms after the onset of the outcome stimulus. Larger positive values correspond to a greater MFN effect. The MFN is indicated by the focus of activity at the Fz electrode (designated by the arrow). The best-fitting dipole model of the generator of the MFN is shown as a red sphere centered in the ACC on a canonical magnetic resonance imaging template of the human head (9).


Gehring and Willoughby, 2002 *Science*

## Error, or motivation?




Gehring and Willoughby, 2002 *Science*

Effect may depend on *relevant* dimension of feedback



Nieuwenhuis, Yeung, Holroyd, Schurger, & Cohen (2004), *Cerebral Cortex*

## FRN may be absence of Reward Positivity




Foti et al. (2011), *HBM*

## Black Jack Study

- 20 Problem Gamblers, 20 Controls
- Black Jack


## FRN and Problem Gambling

Why do Gamblers Gamble?



Hewig et al. (2010). *Biological Psychiatry*

## Black Jack Study



Prob "hit" at 16

Hewig et al. (2010). *Biological Psychiatry*