Advanced Signal Processing II (aka Acronym Day)

PCA ICA Removal of OCULAR artifacts with ICA (and lots of acronyms) BESA Simultaneous EEG with ICA and fMRI!

nternational Trans Day of Visibility, International Trans Day of Visibility,

Announcements 5/5/25

Paper/Proposal due date: May 7, 11:59 pm

- \triangleright Be sure to review the <u>Guidelines</u> (On course webpage, Link in D2L)
- > Pay special attention to methods
 - ≻ Tasks
 - Recording parameters
 - Signal processing to obtain metrics
 - > Analysis plan related to hypotheses
- Take-home exam available tonight on D2L, due May 12 (11:59 pm) (Announcement will alert you)
- Student Course Surveys complete by last day of semester (May 7)

≻501B Lab Section

- May 5 End of Day
 ERP Analysis done (Email Kelly)
- ➢ May 9 End of Day
 ► ERP Report Due

Course Evaluations

- ➢ Your opinions are ...
 - ➢Anonymous
 - ➤Valued
 - >Important to create course improvements
- Find the link:
 - ►D2L
 - Emails that pester you
 - >Or directly: tceonline.oia.arizona.edu

Advanced Signal Processing II (aka Acronym Day)

PCA ICA Removal of OCULAR artifacts with ICA (and lots of acronyms) BESA Simultaneous EEG with ICA and fMRI!

Advanced Signal Processing II (aka Acronym Day)

But first, two things we did not Removal of OCULAR artifacts with ICA (and lots of acronyms) get to last Alme... Simultaneous EEG with ICA and fMRI!

The Problem of Latency Jitter

The averaging assumption of invariance in signal is not always warranted

- Especially for the later endogenous components
- ➢ To the extent that the signal varies from trial to trial, the average will produce potentially misleading results
- > Two common possibilities:
 - Smearing of components;
 - will underestimate amplitude of component (especially a problem if comparing groups, one group with more latency jitter)
 - Bimodal or multi-bumped components

A No r	No noise	
######	######	
#RIGHT	######	
## # # # #	##LEFT	
######	######	
a b		
Noise		
NRIGHT	KWSMNT	
BMJUKM	UYRMUD	
EQĖIKM	VTFMZS	
KEHEHG	ILEFTA	
c <u>1</u> ° d		



The Solution

The Woody Adaptive Filter (Woody, 1967)



The Solution

- The Woody Adaptive Filter (Woody, 1967)
- Based on Cross-correlation
 - Assumptions less restrictive than averaging methods
 - ≻Waveform (morphology) must be constant across trials
 - Latency need not be constant

Details

- Cross-correlational series
 - For two waveforms the correlation between each of them is computed
 - first with no lag in time
 a1, a2, ..., an
 b1, b2, ... bn
 - then with one lagged with respect to the other a1, a2, ..., an-1 b2, b3, ... bn
 - A series of correlation values is obtained by progressively increasing the size of the lag

The Basic Idea

Sine



The Basic Idea



Sine

Cosine

Cross-Correlation

The Basic Idea



See ... CrossCorr_Sin_Cos.m

Cosine

Sine

Cross-Correlation

More Details

- Can be used as a "template matching" procedure
- Compare running average with raw EEG epochs
- This is a method of single-trial signal detection:
 - First create a template: either predetermined (e.g., sine wave) or empirically determined (e.g., average)
 - Then calculate cross-correlational series between each raw EEG epoch and the template
 - ➢ If some maximum correlation achieved, conclude signal is present
 - ➢ If correlation not achieved conclude absent
 - ➤ This can also be used as a method of determining the latency of a component (by examining the trial-by-trial shifts), or of determining the variability in response for a given individual (again by examining the trial-by-trail shifts)

Woody's Instantiation

- The Woody Adaptive Filter (Charles Woody, 1967) is a special case and application of cross correlational technique
- The term "adaptive" refers to the fact that the template is not established a priori, but generated and updated by an iterative procedure from the data themselves
- Procedure
 - Initial template is usually either a half cycle of a sine or triangle wave, or the unfiltered average of single trials
 - Cross-lagged correlations (or sometimes covariances) are then computed between each trial and this template typically over a limited range of samples (e.g., region of P300, not over "invariant" components)
 - Each trial is then shifted to align it with the template at the value which yields the maximum cross correlation (or covariance)
 - A new template is then generated by averaging together these time-shifted epochs
 - Procedure is repeated using this new average as the template
 - Repeated until the maximal values of the cross correlation become stable
 - \triangleright Often, average cross-correlation value increment monitored; if <u>r</u> increases < .005 or .001, then stability achieved
- Some implementations, trials which do not reach a minimum criterion (e.g., .30-.50) are discarded from subsequent template construction and perhaps from subsequent analysis altogether

Woody Filtering Demo!





Validity

- Seems to do a fair job of improving signal extraction if a few iterations are used and if the original signal itself is singly peaked
- Wastell(1977) reports a decline in the validity of the procedure if numerous iterations are used
- Therefore, unlike averaging, Woody filtering can only improve signal-to-noise ratio over a definite limit
- Suggests also that Woody may not be the solution under conditions of very low signal-to-noise ratio

Advanced Signal Processing II (aka Acronym Day)

But first, two things we did not get to last time...

Using Scalp Topography to Infer Different Generators

- Assumption is that if there are different source generators between, there will be different resultant scalp distributions
- Therefore would expect to find a Scalp site by Condition interaction in ANOVA
- The Problem (Wood & McCarthy, 1985)
 - Potentials do not propagate to scalp in strictly additive manner
 - Same source at different strengths can produce a Scalp site by Condition interaction







The Solution

>Normalization

- ➢ For each condition, scale data (e.g. by dividing by site of maximum amplitude)
- Eliminates any overall condition main effect
 - Condition main effect must be assessed in standard (non-scaled) ANOVA
 - Scaled data now lead to an interpretable interaction
 - ➢ If interaction survives scaling, then one can reasonably infer different intra-cranial generators

A New Problem

- Urbach & Kutas (2002) point out that the solution is not a solution! It's intractable
 - For single point source that is invariant in rotation, perhaps Wood & McCarthy were right
 - ➢But when dipole rotates (e.g. on a gyrus), changes polarity, the W&M strategy will not work
 - When there are multiple generators, with changes in *relative* strength, W&M strategy will not work



If, and only if...

- ➢ W&M procedure produces valid inferences if and only if two generator distributions G1, G2, are multiplicatively related
- > Two generator distributions are multiplicatively related iff:
 - \geq 1. The locations of the generators are all the same AND
 - \geq 2. The polarities of the generators are all the same AND
 - ➤ 3. The intensities of the generators differ in overall strength, not *relative* strength
- But how would you ever know, unless you knew where the generators were
 - \succ ... in which case you would not be using the W&M procedure!

So, where's that leave us?

If you scale the amplitudes and there is no interaction between condition and site, then the generators are not different

≻But if there is such an interaction, you don't know whether:

generators differ in location OR

- generators differ in polarity OR
- > generators differ in *relative* strength
- So a nonsignificant effect is informative

Advanced Signal Processing II (aka Acronym Day)

PCA ICA Removal of OCULAR artifacts with ICA (and lots of acronyms) BESA Simultaneous EEG with ICA and fMRI!







Dimensionality explosions!

32, 64, 128, 256!!!



Principal Components Analysis

A method for reducing massive data setsSee Handout for gory details

PCA (1): The Data matrix



- Data Matrix above shows only one site could have multiple sites by adding rows for each subject
- This data matrix is for "temporal PCA" but one have channels by time points matrix and transpose for "spatial PCA"

PCA (2): The Score matrix

S _{Nxm} =	[s ₁ , s ₂ , s ₃ ,, s _m	Where N = Number subjects
Subject #1	s ₁ , s ₂ , s ₃ ,, s _m	m = Number of components
Subject #2	s ₁ , s ₂ , s ₃ ,, s _m	s = score on
Subject #3		component 1, 2,
 Subject #N	 s ₁ , s ₂ , s ₃ ,, s _m]	

- These scores for each subject are optimally weighted composites of the original data, designed to capture as much variance as possible with as few scores as possible.
- ▶ But for conceptual ease, imagine 5 scores: P1, N1, P2, N2, P3 amplitude

PCA (3): The Loading matrix (to guess what components mean)



(100-Hz digitizing rate) in the waveforms.

Spatial PCA on Sample Data



PCA version

PCA (3b): The Loading Map (for Spatial PCA)


Reminder: The ERP from which it derives



PCA Component 2



PCA (4): Reconstructing Data Matrix

 $> \mathbf{D}_{Nxn} \sim = \mathbf{S}_{Nxm} * \mathbf{L}_{mxn}$

This reconstructed Data matrix will differ slightly from the original Data matrix because not all n components are used.

➤ To the extent that the m components account for most of the variance in the original data set, the reconstructed data matrix will closely approximate the original data matrix.

PCA (4): Caveat Emptor

- PCA is a linear model; assumes the components sum together without interaction to produce the actual waveform
- Sources of variance are orthogonal; if two sources are highly correlated, may result in a composite PCA component reflecting both
- Component invariability in terms of latency jitter across subjects
 - PCA does not distinguish between variations in amplitude vs variations in latency
 - Especially a problem in comparing control vs pathological groups; pathological groups will typically be more variable
 - > Allen & Collins unpublished simulation study:
 - Two groups: Control & Pathological
 - > Identical waveforms for each group differed only in latency
 - > The two groups differed significantly on three of four principal component scores
 - In other words, if one indiscriminately interprets these as amplitude or morphology differences, one would be WRONG!!!

ICA ... a "better" PCA?

PCA finds orthogonal components

- First PC accounts for most variance
- > Next PC accounts for most remaining variance
- Components will have orthogonal scalp distributions

ICA separates temporally independent components

- > Also known as blind source separation
- May or may not correspond to brain "hotspots" but do represent functional brain networks

≻ See:

http://arnauddelorme.com/ica_for_dummies/

ICA Decomposition



ICA vs PCA



From Tzyy-Ping Jung , presented at EEGLab Workshop, Nov 8,2007

EEG data are mixtures of source signals



Cocktail Party



From Tzyy-Ping Jung , presented at EEGLab Workshop, Nov 8,2007





Gustafson, Allen, Yeh, May (2011), Early Human Development

ICA/EEG Assumptions

- Mixing is linear at electrodes
- Propagation delays are negligible
- Component time courses are independent
- Number of components < number of channels.

From Tzyy-Ping Jung , presented at EEGLab Workshop, Nov 8,2007

27

ICA: The Projection Map



ICA: The Projection Map



ICA: Trial by Trial IC Projection to Pz



PCA Component 2



ICs as Artifacts!

"Clinical" vs Actuarial Approaches

Clinical Versus Actuarial Judgment

ROBYN M. DAWES, DAVID FAUST, PAUL E. MEEHL

Professionals are frequently consulted to diagnose and predict human behavior; optimal treatment and planning often hinge on the consultant's judgmental accuracy. The consultant may rely on one of two contrasting approaches to decision-making—the clinical and actuarial methods. Research comparing these two approaches shows the actuarial method to be superior. Factors underlying the greater accuracy of actuarial methods, sources of resistance to the scientific findings, and the benefits of increased reliance on actuarial approaches are discussed.

a clinical practitioner. A clinician in psychiatry or medicine may use the clinical or actuarial method. Conversely, the actuarial method should not be equated with automated decision rules alone. For example, computers can automate clinical judgments. The computer can be programmed to yield the description "dependency traits," just as the clinical judge would, whenever a certain response appears on a psychological test. To be truly actuarial, interpretations must be both automatic (that is, prespecified or routinized) and based on empirically established relations.

Virtually any type of data is amenable to actuarial interpretation. For example, interview observations can be coded quantitatively (patient appears withdrawn: [1] yes, [2] no). It is thereby possible to incorporate qualitative observations and quantitative data into

Dawes, R.M., Faust, D., & Meehl, P.E.(1989). Science, 243, 1668-1674.

"Clinical" vs Actuarial Approaches

≻Human raters

Good source of possible algorithms

- ► Lousy at reliably implementing them
 - ≻Inter-rater

≻Intra-rater

>Actuarial methods

>Always arrive at the same conclusion

>Weight variables according to *actual* predictive power

Dawes, R.M., Faust, D., & Meehl, P.E.(1989). Science, 243, 1668-1674.

ICs as Artifacts!

ADJUST: An automatic EEG artifact detector based on the joint use of spatial and temporal features

ICs as Artifacts!

MARA (Multiple Artifact Rejection Algorithm) FASTER (Fully Automated Statistical Thresholding for EEG artifact Rejection) SASICA (a tool for implementing these and more)...

Tool	Artifact type	Measure	Abbreviation
SASICA	Blinks/vertical eye movements	Correlation with vertical EOG electrodes	CorrV
	Horizontal eye movements	Correlation with horizontal EOG electrodes	CorrH
	Muscle	Low autocorrelation of time-course	LoAC or AutoCorr
	Bad channel	Focal channel topography	FocCh
	Rare event	Focal trial activity	FocTr
	Non dipolar component	Residual variance	ResVar
	Bad channel	Correlation with Bad channel	CorrCh
FASTER	Eye blinks/saccades	Correlation with EOG electrodes	EOGcorr
	"Pop-Off"	Spatial Kurtosis	SK
	White noise	Slope of the power spectrum	SpecSl
	White noise	Hurst exponent	HE
	White noise	Median slope of time-course	MedGrad
ADJUST	Eye blinks	Temporal Kurtosis	TK
	Eye blinks	Spatial average difference	SAD
	Eye blinks	Spatial variance difference	SVD
	Vertical Eye Movements	Maximum epoch variance	MEV
	Horizontal Eye Movements	Spatial eye difference	SED
	Generic Discontinuities	Generic discontinuity spatial feature	GDSF

Table 1Measures computed by the three automated tools evaluated here. Abbreviations refer to those used in figures and throughout the paper.



Blink components

Α

. .

Expected properties

Frontal topography

Large amplitude

Opposite polarity below the eyes

> No peak at physiological frequencies

High correlation with vertical EOGs

High eye movement related measures





Horizontal eye movement components

D

Component 13 activity (global offset 0.027) Component 1 activity (global offset -0.067) Е F 700 DS#01 / IC013 DS#03 / IC001 600 600 Expected properties 500 500 400 400 300 300 Opposite sign bilateral frontal topography 200 200 100 σ \mathbf{t} 0.7 .0.1 Step-like 2000 1500 1000 500 0 500 -600 ð. Activity power spectrum Activity power spectrum events -0 -20 Opposite polarity 20 40 around the eyes -40 25 30 35 40 20 25 30 35 10 15 20 45 10 15 -5 Frequency (Hz) Frequency (Hz) No peak at 2*T) 2"Th physiological frequencies SASICA TH SASICA TI LoAC ForCh ForTr LoSNR ResV ConV ConH High correlation with vertical/horizontal EOGs 2*Th 2"Th ADJUST: HEM Th ADJUST: OK Th SAD SED GDSF MEV TK GAB SED High 2*Th 2'Th FASTER: FASTER: eye movement EOO/Correl **EOGCornel** related measures

500 40 45 50 LoAC FeeCh FeeTr LoSNR ReeV CorrH CorrH GDSF MEV TK MedGrad SpecSI SK. HE EOGCorr ModGrad SpecSI SK HE EOGCorr

Non-artifact components may be mistaken for ocular components



Muscle components

А

Expected properties

Focal topography

Steady noisy time courses dissipating / building up across trials

Power at high frequencies

> High noise measures



Bad Channel components

Expected properties

Focal (one channel) topography

Noisy time course

High correlation with marked bad channel

High spatial / intertrial noise measures



Ambiguous mixture components

D



Rare Events



Eye blinks

Features used

- Spatial Average Difference (SAD)
- Temporal Kurtosis (TK)

Frontal distribution

High power in delta frequency band





Vertical Eye Movement

Features used

- Spatial Average Difference (SAD)
- Maximum Epoch Variance (MEV)

Frontal distribution similar to that of an eye blink



Horizontal Eye Movement

Features used

- Spatial Eye Difference (SED)
- Maximum Epoch Variance (MEV)

Frontal distribution in anti-phase (one positive and one negative)



Generic Discontinuities

Features used

- Generic Discontinuities Spatial Feature (GDSF)
- Maximum Epoch Variance (MEV)

Variable distribution

- Sudden amplitude fluctuations with no spatial preference
 - Could be present in as little as one or 2 trials, and limited to 1 channel









After Smith, Reznik, Stewart, Allen (2017)
Neural Sources of EEG

Inverse solution is not unique



0

A single pattern of neural activity will produce a unique scalp map

BUT ... A single scalp map could have been produced by an infinite number of patterns of neural activity

From Tzyy-Ping Jung , presented at EEGLab Workshop, Nov 8,2007

Source Analysis

- BESA -- Brain Electrical Source Analysis
- This is a model-fitting procedure for estimating intracranial sources underlying ERPs
 - Estimate -- if model fits, then data are consistent with these sources; yet there is no unique solution
 - ► Not for ongoing EEG -- too many sources

- Imagine a data matrix of ERPs:
- V_{Cxn} (# Channels by # timepoints)
- Note that this is really the result of the subtraction of the activity at the reference from the activity at the these sites; i.e.,

$$\mathbf{V}_{\mathbf{Cxn}} = \mathbf{U}_{\mathbf{Cxn}} - \mathbf{R}_{\mathbf{Cxn}}$$

Note: the reference matrix has identical rows! Thus BESA Presumes that all channels referenced to the same reference!

Reconstruct a data matrix that includes not only the original channels, but the implicit channel (reference) as well:
 U_{Exn} (# electrodes = # channels+1),

which represents the activity at each electrode with respect to an average reference (i.e., the average of all channels)

Now this matrix U_{Exn} can be decomposed into
 a set of sources: S_{Sxn} (# Sources by # timepoints)
 a set of attenuation coefficients C_{ExS}
 so that U_{Exn} = C_{ExS} S_{Sxn}

The attenuation matrix is determined by:

- \succ the <u>geometry</u> between the source and the electrodes
- ➤ the nature of the conductance of the three-layer head model (Brain, Skull, Scalp);
 - \triangleright the skull is less conductive than the layers on either side
 - \succ this results in a spatial smearing of potentials as they cross the skull
 - ➤ the skull produces the equivalent of a brain that is 60% of the radius of the outer scalp (rather than the "true" figure of ~84%)





Fig. 4. Coronal scalp potential distribution of a radial equivalent dipole modeling activity of superficial cortex. The dipole is oriented inward to mimic, for example, excitatory pyramidal cell activation at the apical dendrites, producing surface negativity. neglecting the shielding effect, i.e. taking an eccentricity of about 80% in a homogeneous head model, results in a narrow focus, similar to the epicortically recorded topography (top). Adequate reduction of equivalent eccentricity results in a realistic scalp topography, which is much more widespread and exhibits a positive maximum on the opposite side of the sphere (bottom). The simulated waveforms at the vertex (C_z) and at equidistant (20^{*}) electrodes over both hemispheres depict a monophasic activity arising with some delay after stimulus delivery.





Fig. 5. Coronal scalp distribution of a tangential dipole modeling fissural cortical activity. As explained for figure 4, the correctly transformed eccentricity in the homogeneous head model (bottom) results in a realistic scalp topography with widespread positive and negative maxima to either side of the actual location of the source. Note that in the quasistatic approach a single dipole source contributes the same waveform at all electrodes. Only the attenuation factor and the sign vary with electrode site.

- \triangleright Note that the decomposition of U into C and S results in
 - ➤ an electroanatomical time-independent matrix (C) that reflects that anatomical substrates do not move around in the head
 - ➤ a time-variant dipole source potential matrix that represents the change in activity of each source over time



BESA Vs PCA Vs ICA (a battle of acronyms)

- This decomposition is akin to PCA/ICA
 DCA and ICA have sources and propagation coefficients
 - > PCA and ICA have sources and propagation coefficients
 - PCA solutions are constrained by orthogonality of components, and by those that account for greatest common variance
 - ICA constrained to find temporally independent components
 - BESA solutions are constrained by the geometry of the head, the volume conduction of the dipoles, and the anatomical constraints dictated by the user (e.g., inside the head, symmetrical, not in the ventricles, must not be in the brainstem after a certain point in time, etc...)

BESA Vs PCA Vs ICA continued

- Like PCA/ICA, the reconstruction of the original data set will be imperfect
 - With all methods. better chance of reconstructing the original matrix if data are reliable
 - If you capture the important sources, the reconstruction should be very good (i.e., small residual variance)
 - It is useful to attempt to upset a solution by inserting another source and seeing if:

 \succ the original solution is stable

➤ the new source accounts for any substantial variance

Can do dipole localization (BESA) on an IC!

Dipole Fitting PCA ICA

You can try it!



Implementations

- \succ BESA can be used:
 - in a strict hypothesis-testing manner by designating sources a priori and testing the fit
 - in an exploratory/optimizing manner by allowing the program to iteratively minimize the residual variance (between observed and reconstructed waveforms) by:
 - moving dipoles
 - changing the orientation of dipoles
 - altering the time-by-activity function of the dipoles

BESA – Did it work?

- ➢ In the end, the adequacy of your solution will be judged by
 - stability of your solution:
 - against insertion of additional dipoles
 - across multiple subjects
 - anatomical feasibility
 - Follow-up tests with patients with lesions
 - Reviewer 2!

Are you nutz?

EEG IN THE MRI

Recording EEG in fMRI environments: Oodles of Issues

- ► EEG can be bad for fMRI
 - > Wires and electrodes can be ferromagnetic = TROUBLE
 - > Wires and electrodes can be paramagnetic = less trouble
- > MRI and fMRI can be bad for EEG
 - Gradient switching creates huge artifact for EEG
 - Movement in Magnetic fields creates current in any conductive medium (e.g. wires!)
 - High frequency current can make wires HOT and RF is 127.68 MHz at 3T – that's fast, and can create mega-hurts!
 - ≻ Thus in-line 10K resistor

Special Caps



- Need conductive material
 That will not heat up
 That will not pose hazard in strong magnetic field
- That includes inline resistor to prevent any induced current from reaching the subject
- That includes Styrofoam head at no charge





1. Hydrogen protons, positively charged particles in the hydrogen molecule's nucleus, normally spin in random directions



2. Protons wobble in alignment with magnetic fields of varying intensity; frequency of wobble is proportionate to strength of individual magnetic field







- equals the frequency of wobble of certain protons, knocks those protons out ofalignment
- 4. When radio signal ceases, protons snap back into alignment with magnetic field, emitting a radio signal of their own, that announces the presence of a specific tissue

- Faraday's law of induction...
 - induced electromotive force is proportional to the time derivative of the magnetic flux
 - Flux = summation of the magnetic field perpendicular to the circuit plane over the area circuit
 - $\bullet \epsilon = d\Phi/dt$
- Can reflect:
 - changes in the field (gradient
 - Changes in the circuit geomet field due to body motion



ne







RF pulses

- ✤ For 3T = 127.6 MHz
- Amplifier frequency range = DC-3.0 KHz
- Artifacts thus attenuated, but still range overwhelm the EEG signal

Gradient Switching

- Artifact approximates differential waveform of the gradient pulse
- Polarity and amplitude varies across channels

Frequency ≈ 500-900 Hz

- EEG dominated by
 - harmonics of slice repetition frequency (≈10-25 Hz)
 - convolved with harmonics of volume repetition frequency (~0.2-2 Hz)
- Artifacts in range from 1000-10,000 μV!



RF = radiofrequency wave;

- Gs = slice selection gradient
- Gp = phase encoding gradient
- Gr = readout gradient
- a = Fat suppression pulses (1-3-3-1 pulses)
- b = slice selection RF
- c, d, h = spoilers
- e = slice selection gradient
- f = dephasing and rephasing gradient
- g = readout gradient
- = EEG artifact corresponding to letter

Ritter et al., 2009

Average Artifact (across 1 TR)



Average Artifact (0-60 msec)



Artifact (across several TRs)



Faraday's law of induction...

- induced electromotive force is proportional to the time derivative of the magnetic flux
- Flux = summation of the magnetic field perpendicular to the circuit plane over the area circuit
- $\bullet \epsilon = d\Phi/dt$

Can reflect:

- changes in the field (gradient switching, RF)
- Changes in the circuit geometry or position relative to the field due to body motion



- ✤ Two types of movement:
 - Axial nodding
 - Expansion at lateral sites
- Motion of blood (flow) can lead to "Hall effect"
- Voltage difference on opposite sides of a moving conductor through which current is flowing, when within a strong magnetic field
- Note field-strength dependent nature of the artifact

Debener et al., 2008

EEG in Magnet (no scanning)



Simulated EKG Artifact







Axial rotation - low frequency spatiallydistributed effect, with polarity reversal

Lateral balloon expansion - locally circumscribed artifact

Debener et al., 2009

Ohmagawd... Help me in

REMOVING THOSE PESKY ARTIFACTS!
Gradiant/RF removal via moving average subtraction



FASTR: FMRI Artifact Slice Template Removal

- Part of FMRIB Plug-in for EEGLAB
- Upsample to at least 20K Hz
- Align slices for slight jitter in timing
- Moving Window approach with subtraction
- PCA on artifact residuals form Optimum Basis Set (OBS) to reduce residual artifacts by 90%
- Downsample to original rate
- Sample Results.....

Before



After



Alternatively ... BrainVision

- Sync EEG clock to MR clock
- No jitter in timing, no need to upsample (recorded at 5000 hz)
- Moving Window approach with subtraction
- Downsample to original rate
- Sample Results.....





ECG-related removal via moving average subtraction (Allen et al. 1998)



Fig. 5 Schematic of the average artefact subtraction procedure. For each channel, a waveform template is generated by averaging EEG epochs over adjacent cardiac cycles, with the time- locking event being derived from the ECG. The template generation is combined with a moving average procedure, and new templates are generated for each cardiac cycle. The procedure is repeated for each EEG channel

There may be residual crud (RC)



There may be residual crud (RC)



Simultaneous EEG and RSfMRI (following ICA!)

Multi-modal Imaging

Create RS-fMRI network with ACC seeds







Allen, Hewig, Miltner, Hecht, & Schnyer, in preparation

EEG Alpha Asymmetry is Negatively Correlated with IFG Connectivity in Two ACC-seeded Resting State Networks

Spatially-enhanced EEG asymmetry (using CSD transform) at sites F8-F7 is related to resting state connectivity between left inferior frontal gyrus and two ACC-seeded networks.



Allen, Hewig, Miltner, Hecht, & Schnyer, in preparation

EEG-fMRI Synopsis

- Less relative left frontal activity (indexed by EEG) is related to increased connectivity of left IFG to two ACC-seeded RS networks
- Consistent with:
 - Hyper-connectivity in RSfMRI emotion networks in MDD (e.g., Grecius et al., 2007; Sheline et al., 2010)
 - Frontal EEG asymmetry findings of less relative left frontal activity in risk for MDD.
- Alpha power may regulate network connectivity
 - Note: Between vs Within Subjects

BETWEEN-SUBJECTS' DATA DOES NOT NECESSARILY SUPPORT A WITHIN-SUBJECTS' INTERPRETATION

Within Subjects' Moderation of RSfMRI Connectivity

- Calculate F8-F7 alpha asymmetry for each TR
 - EEG leads TR by 4.096 seconds
- Median split into high (left) and low (right)
- Entered as moderator in PPI approach (cf. Friston et al., 1997)
 - Tests whether strength of connectivity to seed region varies as a function of the moderator

Within Subjects' Moderation of RSfMRI Connectivity





Dorsal ACC Seed

Greater Connectivity with Less Left Frontal Alpha or Greater Left Frontal Alpha

Allen, Hewig, Miltner, Hecht, & Schnyer, in preparation

Within (red) and Between (blue) Within-subject effects more extensive



Cognitive Control over Emotion

IFG has a key role in mediating the success of cognitive control over emotional stimuli

Cognitive Control over Emotion

- ✦ Left IFG: Language and self-referential Ġ NO ONTAI processing FRONTR MED. SUBFRONT 0 ERFRONTAL Ц Ω PARAMESIAL F. ш ЕR Σ ۵. ENTRAL IPRECENTRAL, S ENTRAL A ROST CEN NFLECTED F CCD. CENTRE Ģ RAL POSTCENT PAR FIRATI Q 00000 CCIPITAL PAROCON
- Right IFG: Attentional control
 - behavioral inhibition
 - suppression of unwanted thoughts
 - attention shifting
 - efforts to reappraise
 emotional stimuli

Cognitive Control over Emotion





Psychophysiology -- Synopsis

- Psychophysiology is inherently interdisciplinary, and systemic
- Principles learned here can apply to a wide range of physiological signals
 - ►Recording
 - ➢Processing
 - >Interpretation

Psychophysiology -- Synopsis

- Ultimately we obtain correlates of behavior and experience
 - Psychophysiological Correlates are not privileged; they are no better, no worse, than any other correlate of behavior and experience
- The utility of these correlates like any correlates in science hinges upon:
 - ➢ good experimental design
 - \succ strong theoretically driven hypothesis testing
 - ➤ the development of a nomological net, a set of interrelationships among tangible measures and constructs that place the findings in a larger theoretical context, and lend construct validity to the measures and findings